

ANTI- AMYLOID ANTIBODIES, COMPOSITIONS, METHODS AND USES

This application claims priority to Provisional Applications S/N 60/458,474;
S/N 60/458,469; S/N 60/458,510; and S/N 60/458,509, all filed March 28, 2003, and are
5 entirely incorporated herein by reference.

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

The present invention relates to antibodies, including specified portions or variants,
10 specific for at least one beta-amyloid (amyloid) protein or fragment thereof, as well as anti-
idiotype antibodies, and nucleic acids encoding such anti-amyloid antibodies, complementary
nucleic acids, vectors, host cells, and methods of making and using thereof, including therapeutic
formulations, administration and devices.

RELATED ART

Alzheimer's Disease (AD) is a degenerative brain disorder characterized clinically by
progressive loss of memory, cognition, reasoning, judgment and emotional stability that
gradually leads to profound mental deterioration and ultimately death. AD is a very common
cause of progressive mental failure (dementia) in aged humans and is believed to represent the
20 fourth most common medical cause of death in the United States. AD has been observed in races
and ethnic groups worldwide and presents a major present and future public health problem. The
disease is currently estimated to affect about two to three million individuals in the United States
alone. AD is at present incurable. No treatment that effectively prevents AD or reverses its
symptoms and course is currently known

25 The brains of individuals with AD exhibit characteristic lesions termed senile (or
amyloid) plaques, amyloid angiopathy (amyloid deposits in blood vessels) and neurofibrillary
tangles. Large numbers of these lesions, particularly amyloid plaques and neurofibrillary tangles,
are generally found in several areas of the human brain important for memory and cognitive
function in patients with AD. Smaller numbers of these lesions in a more restricted anatomical
30 distribution are also found in the brains of most aged humans who do not have clinical AD.
Amyloid plaques and amyloid angiopathy also characterize the brains of individuals with
Trisomy 21 (Down's Syndrome), Diffuse Lewy Body Disease and Hereditary Cerebral
Hemorrhage with Amyloidosis of the Dutch-Type (HCHWA-D).

A major constituent of amyloid plaques are a variety amyloid-beta (A β) peptides which
35 are produced by cleavage of the β -amyloid precursor protein (APP). While in the past

there was significant scientific debate over whether the plaques and tangles are a cause or are merely the result of Alzheimer's disease, recent discoveries indicate that amyloid plaque is a causative precursor or factor. In particular, it has been discovered that the production of A β peptides can result from mutations in the gene encoding amyloid precursor protein, a protein which when normally processed will not produce the A β peptides. The identification of mutations in the amyloid precursor protein gene which cause familial, early onset Alzheimer's disease is the strongest evidence that amyloid metabolism is the central event in the pathogenic process underlying the disease. It is presently believed that a normal (non-pathogenic) processing of the APP protein occurs via cleavage by an "alpha-secretase" which cleaves between amino acids 16 and 17 of the A β peptide region within the protein. It is further believed that pathogenic processing occurs in part via "beta-secretases" which cleave at the amino-terminus of the A β peptide region within the precursor protein. Beta amyloid protein is also thought to be potentially associated with other neurological and some cardiovascular disorders.

Accordingly, there is a need to provide anti-amyloid antibodies or fragments that overcome one more of these problems, as well as improvements over known antibodies or fragments thereof.

SUMMARY OF THE INVENTION

The present invention provides isolated human, primate, rodent, mammalian, chimeric, humanized and/or CDR-grafted anti-amyloid antibodies, immunoglobulins, fragments, cleavage products and other specified portions and variants thereof, as well as anti-amyloid antibody compositions, encoding or complementary nucleic acids, vectors, host cells, compositions, formulations, devices, transgenic animals, transgenic plants, and methods of making and using thereof, as described and enabled herein, in combination with what is known in the art.

The present invention also provides at least one isolated anti-amyloid antibody as described herein. An antibody according to the present invention includes any protein or peptide containing molecule that comprises at least a portion of an immunoglobulin molecule, such as but not limited to, at least one ligand binding portion (LBP), such as but not limited to, a complementarity determining region (CDR) of a heavy or light chain (e.g., comprising at least one of SEQ ID NOS:42-47, 53-58, 63-68, 73-78) or a ligand binding portion thereof, a heavy chain or light chain variable region (e.g., comprising at least one of 10-125 contiguous amino acids of at least one of SEQ ID NOS:1-30, or at least one FR1, FR2, FR3, FR4 or fragment thereof as described in Table 1, further optionally comprising at least one substitution,

insertion or deletion as provided in Figures 1-41), a heavy chain or light chain constant region (e.g., comprising at least one of 10-384 contiguous amino acids of at least one of SEQ ID NOS:31-41, or at least one CH1, hinge1, hinge2, hinge 3, hinge4, CH2, CH3 or fragment thereof as described in Table 1, further optionally comprising at least one substitution, insertion or deletion as provided in Figures 1-41), a framework region, or any portion thereof, that can be incorporated into an antibody of the present invention. An antibody of the invention can include or be derived from any mammal, such as but not limited to a human, a mouse, a rabbit, a rat, a rodent, a primate, or any combination thereof, and the like.

The present invention provides, in one aspect, isolated nucleic acid molecules comprising, complementary, or hybridizing to, a polynucleotide encoding specific anti-amyloid antibodies, comprising at least one specified sequence, domain, portion or variant thereof. The present invention further provides recombinant vectors comprising said anti-amyloid antibody nucleic acid molecules, host cells containing such nucleic acids and/or recombinant vectors, as well as methods of making and/or using such antibody nucleic acids, vectors and/or host cells.

The present invention also provides at least one anti-amyloid antibody or specified portion or variant, comprising at least one amyloid binding sequence and at least 10-384 contiguous amino acids of at least one of SEQ ID NOS:1-41, or at least one FR1, FR2, FR3, FR4, CH1, hinge1, hinge2, hinge 3, hinge4, CH2, CH3 or fragment thereof as described in Table 1, further optionally comprising at least one substitution, insertion or deletion as provided in Figures 1-41, or as known in the art.

At least one antibody of the invention binds at least one specified epitope specific to at least one amyloid protein, subunit, fragment, portion or any combination thereof. The at least one epitope can comprise at least one antibody binding region that comprises at least one portion of said protein, which epitope is preferably comprised of at least 1-5 amino acids of at least one portion thereof, such as but not limited to, at least one functional, extracellular, soluble, hydrophilic, external or cytoplasmic domain of said protein, or any portion thereof.

The at least one antibody can optionally comprise at least one specified portion of at least one complementarity determining region (CDR) (e.g., CDR1, CDR2 or CDR3 of the heavy or light chain variable region) and optionally further comprising at least one constant or variable framework region or any portion thereof. The at least one antibody amino acid sequence can further optionally comprise at least one specified substitution, insertion or deletion as described herein or as known in the art.

The present invention also provides at least one isolated anti-amyloid antibody as described herein, wherein the antibody has at least one activity, such as, but not limited to one known amyloid protein assay. An anti-amyloid antibody can thus be screened for a

corresponding activity according to known methods, such as but not limited to, at least one biological activity towards an amyloid protein.

5 The present invention further provides at least one amyloid anti-idiotypic antibody to at least one amyloid antibody of the present invention. The anti-idiotypic antibody includes any protein or peptide containing molecule that comprises at least a portion of an immunoglobulin molecule, such as but not limited to at least one ligand binding portion (LBP), such as but not limited to a complementarity determining region (CDR) of a heavy or light chain, or a ligand binding portion thereof, a heavy chain or light chain variable region, a heavy chain or light chain constant region, a framework region, or any portion thereof, that can be incorporated into an antibody of the present invention. An antibody of the invention can include or be derived from any mammal, such as but not limited to a human, a mouse, a rabbit, a rat, a rodent, a primate, and the like.

15 The present invention provides, in one aspect, isolated nucleic acid molecules comprising, complementary, or hybridizing to, a polynucleotide encoding at least one amyloid anti-idiotypic antibody, comprising at least one specified sequence, domain, portion or variant thereof. The present invention further provides recombinant vectors comprising said amyloid anti-idiotypic antibody encoding nucleic acid molecules, host cells containing such nucleic acids and/or recombinant vectors, as well as methods of making and/or using such anti-idiotypic antibody nucleic acids, vectors and/or host cells.

20 The present invention also provides at least one method for expressing at least one anti-amyloid antibody, or amyloid anti-idiotypic antibody, in a host cell, comprising culturing a host cell as described herein under conditions wherein at least one anti-amyloid antibody is expressed in detectable and/or recoverable amounts.

25 The present invention also provides at least one composition comprising (a) an isolated anti-amyloid antibody encoding nucleic acid and/or antibody as described herein; and (b) a suitable carrier or diluent. The carrier or diluent can optionally be pharmaceutically acceptable, according to known carriers or diluents. The composition can optionally further comprise at least one further compound, protein or composition.

30 The present invention further provides at least one anti-amyloid antibody method or composition, for administering a therapeutically effective amount to modulate or treat at least one amyloid related condition in a cell, tissue, organ, animal or patient and/or, prior to, subsequent to, or during a related condition, as known in the art and/or as described herein.

35 The present invention also provides at least one composition, device and/or method of delivery of a therapeutically or prophylactically effective amount of at least one anti-amyloid antibody, according to the present invention.

The present invention further provides at least one anti-amyloid antibody method or composition, for diagnosing at least one amyloid related condition in a cell, tissue, organ, animal or patient and/or, prior to, subsequent to, or during a related condition, as known in the art and/or as described herein.

5 The present invention also provides at least one composition, device and/or method of delivery for diagnosing of at least one anti-amyloid antibody, according to the present invention.

In one aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising at least one variable region comprising SEQ ID NO:48 or 49.

10 In another aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising either (i) all of the heavy chain complementarity determining regions (CDR) amino acid sequences of SEQ ID NOS:42-44; or (ii) all of the light chain CDR amino acids sequences of SEQ ID NOS:45-47.

15 In another aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising at least one heavy chain or light chain CDR having the amino acid sequence of at least one of SEQ ID NOS: 42-47.

In one aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising at least one variable region comprising SEQ ID NO:59 or 60.

20 In another aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising either (i) all of the heavy chain complementarity determining regions (CDR) amino acid sequences of SEQ ID NOS:53-55; or (ii) all of the light chain CDR amino acids sequences of SEQ ID NOS:56-58.

25 In another aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising at least one heavy chain or light chain CDR having the amino acid sequence of at least one of SEQ ID NOS:53-58.

In one aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising at least one variable region comprising SEQ ID NO:69 or 70.

30 In another aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising either (i) all of the heavy chain complementarity determining regions (CDR) amino acid sequences of SEQ ID NOS:63-65; or (ii) all of the light chain CDR amino acids sequences of SEQ ID NOS:66-68.

In another aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising at least one heavy chain or light chain CDR having the amino acid sequence of at least one of SEQ ID NOS:63-68.

In one aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising at least one variable region comprising SEQ ID NO:79 or 80.

In another aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising either (i) all of the heavy chain complementarity
5 determining regions (CDR) amino acid sequences of SEQ ID NOS:73-75; or (ii) all of the light chain CDR amino acids sequences of SEQ ID NOS:76-78.

In another aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising at least one heavy chain or light chain CDR having the amino acid sequence of at least one of SEQ ID NOS:73-78.

10 In one aspect, the present invention provides at least one isolated mammalian anti-amyloid antibody, comprising at least one human CDR, wherein the antibody specifically binds at least one epitope selected from amino acids 2-7, 3-8, 33-42, or 34-40 of SEQ ID NO:50.

In another aspect, the present invention provides at least one isolated mammalian anti-
15 amyloid antibody, comprising at least one human CDR, wherein the antibody specifically binds at least one epitope comprising at least 1-3, to the entire amino acid sequence of SEQ ID NO:50.

The at least one antibody can optionally further comprise at least one characteristic selected from: (i) bind amyloid with an affinity of at least one selected from at least 10^{-9} M, at
20 least 10^{-10} M, at least 10^{-11} M, or at least 10^{-12} M; and/or (ii) substantially neutralize at least one activity of at least one amyloid protein. Also provided is an isolated nucleic acid encoding at least one isolated mammalian anti-amyloid antibody; an isolated nucleic acid vector comprising the isolated nucleic acid, and/or a prokaryotic or eukaryotic host cell comprising the isolated nucleic acid. The host cell can optionally be at least one selected from COS-1,
25 COS-7, HEK293, BHK21, CHO, BSC-1, Hep G2, 653, SP2/0, 293, HeLa, myeloma, or lymphoma cells, or any derivative, immortalized or transformed cell thereof. Also provided is a method for producing at least one anti-amyloid antibody, comprising translating the antibody encoding nucleic acid under conditions in vitro, in vivo or in situ, such that the amyloid antibody is expressed in detectable or recoverable amounts.

30 Also provided is a composition comprising at least one isolated mammalian anti-amyloid antibody and at least one pharmaceutically acceptable carrier or diluent. The composition can optionally further comprise an effective amount of at least one compound or protein selected from at least one of a detectable label or reporter, an anti-infective drug, a cardiovascular (CV) system drug, a central nervous system (CNS) drug, an autonomic nervous

system (ANS) drug, a respiratory tract drug, a gastrointestinal (GI) tract drug, a hormonal drug, a drug for fluid or electrolyte balance, a hematologic drug, an antineoplastic, an immunomodulation drug, an ophthalmic, otic or nasal drug, a topical drug, a nutritional drug or the like, a TNF antagonist, an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NTHE), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, an erythropoietin, an immunization, an immunoglobulin, an immunosuppressive, a growth hormone, a hormone replacement drug, a radiopharmaceutical, an antidepressant, an antipsychotic, a stimulant, an asthma medication, a beta agonist, an inhaled steroid, an epinephrine or analog, a cytokine, or a cytokine antagonist.

The present invention further provides an anti-idiotypic antibody or fragment that specifically binds at least one isolated mammalian anti-amyloid antibody of the present invention.

Also provided is a method for diagnosing or treating an amyloid related condition in a cell, tissue, organ or animal, comprising

(a) contacting or administering a composition comprising an effective amount of at least one isolated mammalian anti-amyloid antibody of the invention with, or to, the cell, tissue, organ or animal. The method can optionally further comprise using an effective amount of 0.001-50 mg/kilogram per: 1-24 hours, 1-7 days, 1-52 weeks, 1-24 months, 1-30 years (or any range or value therein), of the cells, tissue, organ or animal. The method can optionally further comprise using the contacting or the administering by at least one mode selected from parenteral, subcutaneous, intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelical, intracelebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, intralesional, bolus, vaginal, rectal, buccal, sublingual, intranasal, or transdermal. The method can optionally further comprise administering, prior, concurrently or after the (a) contacting or administering, at least one composition comprising an effective amount of at least one compound or protein selected from at least one of an anti-infective drug, a cardiovascular (CV) system drug, a central nervous system (CNS) drug, an autonomic nervous system (ANS) drug, a respiratory tract drug, a gastrointestinal (GI) tract drug, a hormonal drug, a drug for fluid or electrolyte balance, a hematologic drug, an antineoplastic, an immunomodulation drug, an ophthalmic, otic or nasal drug, a topical drug, a nutritional drug or the like. The method can optionally further comprise administering, prior, concurrently or

after the (a) contacting or administering, at least one composition comprising an effective amount of at least one compound or protein selected from at least one of a detectable label or reporter, a TNF antagonist, an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NTHE), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, an erythropoietin, an immunization, an immunoglobulin, an immunosuppressive, a growth hormone, a hormone replacement drug, a radiopharmaceutical, an antidepressant, an antipsychotic, a stimulant, an asthma medication, a beta agonist, an inhaled steroid, an epinephrine or analog, a cytokine, or a cytokine antagonist.

Also provided is a medical device, comprising at least one isolated mammalian anti-amyloid antibody of the invention, wherein the device is suitable to contacting or administering the at least one anti-amyloid antibody by at least one mode selected from parenteral, subcutaneous, intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelial, intracelebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, intralesional, bolus, vaginal, rectal, buccal, sublingual, intranasal, or transdermal.

Also provided is an article of manufacture for human pharmaceutical or diagnostic use, comprising packaging material and a container comprising a solution or a lyophilized form of at least one isolated mammalian anti-amyloid antibody of the present invention. The article of manufacture can optionally comprise having the container as a component of a parenteral, subcutaneous, intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelial, intracelebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, intralesional, bolus, vaginal, rectal, buccal, sublingual, intranasal, or transdermal delivery device or system.

Also provided is a method for producing at least one isolated mammalian anti-amyloid antibody of the present invention, comprising providing a host cell or transgenic animal or transgenic plant or plant cell capable of expressing in recoverable amounts the antibody. Further provided in the present invention is at least one anti-amyloid antibody produced by the above method.

The present invention further provides any invention described herein.

DESCRIPTION OF THE FIGURES

Figures 1-41 show the heavy/light chain variable/constant region prototype sequences, frameworks/subdomains and substitutions. For each region, multiple sequence alignment of known sequences was performed and a prototype sequence was generated. Framework, CDR and hinge regions are labeled in boxes. Prototype sequence residues are numbered for each amino acid position. A list of amino acid substitutions or gaps (denoted by a "-") observed at each prototype position in the aligned sequences and the number of times of their occurrence are shown below each prototype sequence residue.

Figure 1 depicts Vh1 heavy chain variable region prototype sequences, frameworks and substitutions.

Figure 2 depicts Vh2 heavy chain variable region prototype sequences, frameworks and substitutions.

Figure 3 depicts Vh3a heavy chain variable region prototype sequences, frameworks and substitutions.

Figure 4 depicts Vh3b heavy chain variable region prototype sequences, frameworks and substitutions.

Figure 5 depicts Vh3c heavy chain variable region prototype sequences, frameworks and substitutions.

Figure 6 depicts Vh4 heavy chain variable region prototype sequences, frameworks and substitutions.

Figure 7 depicts Vh5 heavy chain variable region prototype sequences, frameworks and substitutions.

Figure 8 depicts Vh6 heavy chain variable region prototype sequences, frameworks and substitutions.

Figure 9 depicts Vh7 heavy chain variable region prototype sequences, frameworks and substitutions.

Figure 10 depicts $\kappa 1_4$ light chain variable region prototype sequences, frameworks and substitutions.

Figure 11 depicts $\kappa 2$ light chain variable region prototype sequences, frameworks and substitutions.

Figure 12 depicts $\kappa 3$ light chain variable region prototype sequences, frameworks and substitutions.

Figure 13 depicts $\kappa 5$ light chain variable region prototype sequences, frameworks and substitutions.

Figure 14 depicts κ New1 light chain variable region prototype sequences, frameworks and substitutions.

Figure 15 depicts κ New2 light chain variable region prototype sequences, frameworks and substitutions.

5 Figure 16 depicts λ 1a light chain variable region prototype sequences, frameworks and substitutions.

Figure 17 depicts λ 1b light chain variable region prototype sequences, frameworks and substitutions.

10 Figure 18 depicts λ 2 light chain variable region prototype sequences, frameworks and substitutions.

Figure 19 depicts λ 3a light chain variable region prototype sequences, frameworks and substitutions.

Figure 20 depicts λ 3b light chain variable region prototype sequences, frameworks and substitutions.

15 Figure 21 depicts λ 3c light chain variable region prototype sequences, frameworks and substitutions.

Figure 22 depicts λ 3e light chain variable region prototype sequences, frameworks and substitutions.

20 Figure 23 depicts λ 4a light chain variable region prototype sequences, frameworks and substitutions.

Figure 24 depicts λ 4b light chain variable region prototype sequences, frameworks and substitutions.

Figure 25 depicts λ 5 light chain variable region prototype sequences, frameworks and substitutions.

25 Figure 26 depicts λ 6 light chain variable region prototype sequences, frameworks and substitutions.

Figure 27 depicts λ 7 light chain variable region prototype sequences, frameworks and substitutions.

30 Figure 28 depicts λ 8 light chain variable region prototype sequences, frameworks and substitutions.

Figure 29 depicts λ 9 light chain variable region prototype sequences, frameworks and substitutions.

Figure 30 depicts λ 10 light chain variable region prototype sequences, frameworks and substitutions.

Figure 31 depicts IgA1 heavy chain constant region prototype sequences, subdomains and substitutions.

Figure 32 depicts IgA2 heavy chain constant region prototype sequences, subdomains and substitutions.

- 5 Figure 33 depicts IgD heavy chain constant region prototype sequences, subdomains and substitutions.

Figure 34 depicts IgE heavy chain constant region prototype sequences, subdomains and substitutions.

- 10 Figure 35 depicts IgG1 heavy chain constant region prototype sequences, subdomains and substitutions.

Figure 36 depicts IgG2 heavy chain constant region prototype sequences, subdomains and substitutions.

Figure 37 depicts IgG3 heavy chain constant region prototype sequences, subdomains and substitutions.

- 15 Figure 38 depicts IgG4 heavy chain constant region prototype sequences, subdomains and substitutions.

Figure 39 depicts IgM heavy chain constant region prototype sequences, subdomains and substitutions.

Figure 40 depicts Igκ light chain constant region prototype sequences and substitutions.

- 20 Figure 41 depicts Igλ light chain constant region prototype sequences and substitutions.

Figure 42 shows the spot membrane probed by C701 mAb. The human Aβ sequence was scanned with peptides shifted by one amino acid. For sequences of regions:

- 25 1. AEFRHDSGYEVH (SEQ ID NO:83);
 2. EFRHDSGYEVHH (SEQ ID NO:84);
 3. IIGLMVGGVVIA (SEQ ID NO:85);
 4. IGLMVGGVVIA (SEQ ID NO:86);
 5. IGLMVGGVVI (SEQ ID NO:87);
 6. IGLMVGGVV (SEQ ID NO:88);
 7. IGLMVGGV (SEQ ID NO:89);
30 8. IGLMVGG (SEQ ID NO:90);
 9. LMVGGV (SEQ ID NO:91).

Figure 43 shows the spot membrane probed by C705 mAb. The human Aβ sequence was scanned with peptides shifted by one amino acid. For sequences of regions:

- 35 1. DAEFRHDSGYEVHHQ (SEQ ID NO:92);
 2. AEFRHDSGYEVHHQ (SEQ ID NO:93);
 3. DAEFRHDSGYEVH (SEQ ID NO:94);
 4. EFRHDSGYEVHH (SEQ ID NO:95);
 5. EFRHDSGYEVH (SEQ ID NO:96);
 6. DAEFRHDSGY (SEQ ID NO:97);
40 7. DAEFRHDSG (SEQ ID NO:98);
 8. AEFRHDSG (SEQ ID NO:99);
 9. EFRHDSG (SEQ ID NO:100);
 10. EFRHDS (SEQ ID NO:101).

- 45 Figure 44 shows the spot membrane probed by C706 mAb. The human Aβ sequence was scanned with peptides shifted by one amino acid. For sequences of regions:

1. DAEFRHDSGYEVHHQ (SEQ ID NO:102);

2. AEFRHDSGYEVHHQ (SEQ ID NO:103);
 3. AEFRHDSGYEVHH (SEQ ID NO:104);
 4. AEFRHDSGYEVH (SEQ ID NO:105);
 5. DAEFRHDSGYE (SEQ ID NO:106);
 - 5 6. AEFRHDSGYE (SEQ ID NO:107);
 7. DAEFRHDSG (SEQ ID NO:108);
 8. AEFRHDSG (SEQ ID NO:109);
 9. DAEFRHDSG (SEQ ID NO:110);
 10. AEFRHD (SEQ ID NO:111).
- 10 Figure 45 shows the spot membrane probed by C707 mAb. The human A β sequence was scanned with peptides shifted by one amino acid. For sequences of regions:
1. EFRHDSGYEVHHQKL (SEQ ID NO:112);
 2. FRHDSGYEVHHQKL (SEQ ID NO:113);
 3. EVHHQKL VFFAEDV (SEQ ID NO:114);
 - 15 4. YEVHHQKL VFFA (SEQ ID NO:115);
 5. VFFAEDVGSNKGA (SEQ ID NO:116);
 6. KLVFFAEDVGSN (SEQ ID NO:117);
 7. EDVGSNKGAIIG (SEQ ID NO:118);
 8. FFAEDVGSNKG (SEQ ID NO:119);
 - 20 9. SNKGAIIGLMV (SEQ ID NO:120);
 10. FAEDVGSNKG (SEQ ID NO:121);
 11. KGAIIGLMVG (SEQ ID NO:122);
 12. LVFFAEDVG (SEQ ID NO:123);
 13. EDVGSNKG (SEQ ID NO:124);
 - 25 14. KLVFFAED (SEQ ID NO:125);
 15. DVGSNKG (SEQ ID NO:126);
 16. EVHHQKL (SEQ ID NO:127);
 17. QKL VFFA (SEQ ID NO:128);
 18. RHDSGY (SEQ ID NO:129);
 - 30 19. SGYEVH (SEQ ID NO:130);
 20. GVVIAT (SEQ ID NO:131).
- Figure 46 shows the A β ₃₈ binding with anti-A β mAb C705.
- Figure 47A shows the ranking of anti-A β mAbs using a plot of the binding ration of A β ₃₈.
- Figure 47B shows the ranking of anti-A β mAbs using a plot of the binding ration of A β ₄₂.
- 35 Figure 48 shows the effect of anti-A β mAbs on A β ₄₂-induced toxicity in PC12 cells.

DESCRIPTION OF THE INVENTION

The present invention provides isolated, recombinant and/or synthetic anti-amyloid human, primate, rodent, mammalian, chimeric, humanized or CDR-grafted, antibodies and amyloid anti-idiotypic antibodies thereto, as well as compositions and encoding nucleic acid molecules comprising at least one polynucleotide encoding at least one anti-amyloid antibody or anti-idiotypic antibody. The present invention further includes, but is not limited to, methods of making and using such nucleic acids and antibodies and anti-idiotypic antibodies, including diagnostic and therapeutic compositions, methods and devices.

As used herein, an "anti-beta-amyloid antibody," "anti-amyloid antibody," "anti-amyloid antibody portion," or "anti-amyloid antibody fragment" and/or "anti-amyloid antibody variant" and the like include any protein or peptide containing molecule that comprises at least a portion of an immunoglobulin molecule, such as but not limited to at least one complementarity determining region (CDR) of a heavy or light chain or a ligand binding portion thereof, a heavy chain or light chain variable region, a heavy chain or light chain constant region, a framework region, or any portion thereof, or at least one portion of an amyloid receptor or binding protein, which can be incorporated into an antibody of the present invention. Such antibody optionally further affects a specific ligand, such as but not limited to where such antibody modulates, decreases, increases, antagonizes, agonizes, mitigates, alleviates, blocks, inhibits, abrogates and/or interferes with at least one amyloid activity or binding, or with amyloid receptor activity or binding, *in vitro*, *in situ* and/or *in vivo*. As a non-limiting example, a suitable anti-amyloid antibody, specified portion or variant of the present invention can bind at least one amyloid, or specified portions, variants or domains thereof. A suitable anti-amyloid antibody, specified portion, or variant can also optionally affect at least one of amyloid activity or function, such as but not limited to, RNA, DNA or protein synthesis, amyloid release, amyloid receptor signaling, membrane amyloid cleavage, amyloid activity, amyloid production and/or synthesis.

Antibodies can include one or more of at least one CDR, at least one variable region, at least one constant region, at least one heavy chain (e.g., γ_1 , γ_2 , γ_3 , γ_4 , μ , α_1 , α_2 , δ , ϵ), at least one light chain (e.g., κ and λ), or any portion or fragment thereof, and can further comprise interchain and intrachain disulfide bonds, hinge regions, glycosylation sites that can be separated by a hinge region, as well as heavy chains and light chains. Light chains typically have a molecular weight of about 25Kd and heavy chains typically range from 50K-77Kd. Light chains can exist in two distinct forms or isotypes, kappa (κ) and lambda (λ), which can

combine with any of the heavy chain types. All light chains have at least one variable region and at least one constant region. The IgG antibody is considered a typical antibody structure and has two intrachain disulfide bonds in the light chain (one in variable region and one in the constant region), with four in the heavy chain, and such bond encompassing a peptide loop of about 60-70 amino acids comprising a "domain" of about 110 amino acids in the chain. IgG antibodies can be characterized into four classes, IgG1, IgG2, IgG3 and IgG4. Each immunoglobulin class has a different set of functions. The following table summarizes the reported examples of the physicochemical properties of each of the immunoglobulin classes and subclasses.

| Property | IgG1 | IgG2 | IgG3 | IgG4 | IgM | IgA1 | IgA2 | SIgA | IgD | IgE |
|------------------------------|------------|------------|------------|------------|-------|------------|------------|-----------------------|----------|---------|
| Heavy Chain | $\gamma 1$ | $\gamma 1$ | $\gamma 1$ | $\gamma 1$ | μ | $\alpha 1$ | $\alpha 2$ | $\alpha 1 / \alpha 2$ | δ | E |
| Mean Serum conc. (mg/ml) | 9 | 3 | 1 | 0.5 | 1.5 | 3.0 | 0.5 | 0.05 | 0.03 | 0.00005 |
| Sedimentation constant | 7s | 7s | 7s | 7s | 19s | 7s | 7s | 11s | 7s | 8s |
| Mol. Wt. ($\times 10^3$) | 146 | 146 | 170 | 146 | 970 | 160 | 160 | 385 | 184 | 188 |
| Half Life (days) | 5-30 | 5-30 | 2-10 | 5-30 | 5-15 | 2-10 | 2-10 | 1-10 | 1-10 | 1-10 |
| % intravascular distribution | 45 | 45 | 45 | 45 | 80 | 42 | 42 | Trac e | 75 | 50 |
| Carbohydrate (%) | 2-3 | 2-3 | 2-3 | 2-3 | 12 | 7-11 | 7-11 | 7-11 | 9-14 | 12 |

10

The following table summarizes non-limiting examples of antibody effector functions for human antibody classes and subclasses.

| Effector function | IgG1 | IgG2 | IgG3 | IgG4 | IgM | IgA | IgD | IgE |
|---------------------|------|------|------|------|-----|-----|-----|-----|
| Complement fixation | ++ | + | +++ | - | +++ | - | - | - |
| Placental transfer | + | + | + | + | - | - | - | - |
| Binding to Staph A | +++ | +++ | - | +++ | - | - | - | - |
| Binding to Strep G | +++ | +++ | +++ | +++ | - | - | - | - |

Accordingly, the type of antibody or fragment thereof can be selected for use according to the present invention based on the desired characteristics and functions that are desired for a particular therapeutic or diagnostic use, such as but not limited to serum half life, intravascular distribution, complement fixation, etc.

Antibody diversity is generated by at least 5 mechanisms, including (1) the use of multiple genes encoding parts of the antibody; (2) somatic mutation, e.g., primordial V gene mutation during B-cell ontogeny to produce different V genes in different B-cell clones; (3) somatic recombination, e.g., gene segments J1-Jn recombine to join the main part of the V-region gene during B-cell ontogeny; (4) gene conversion where sections of DNA from a number of pseudo V region can be copied into the V region to alter the DNA sequence; and (5) nucleotide addition, e.g., when V and J regions are cut, before joining, and extra nucleotides

may be inserted to code for additional amino acids. Non-limiting examples include, but are not limited to, (i) the selection/recombination of V κ , J, and C κ regions from germ line to B-cell clones to generate kappa chains; (ii) selection/recombination of V λ , J, and C λ regions from germ line to B-cell clones to generate lambda chains; (iii) selection/recombination of V_H, D1-D30 and J_H1-J_H6 genes to form a functional VDJ gene encoding a heavy chain variable region. The above mechanisms work in a coordinated fashion to generate antibody diversity and specificity.

The term "antibody" is further intended to encompass antibodies, digestion fragments, specified portions and variants thereof, including antibody mimetics or comprising portions of antibodies that mimic the structure and/or function of an antibody or specified fragment or portion thereof, including single chain antibodies and fragments thereof. Functional fragments include antigen-binding fragments that bind to a mammalian amyloid. For example, antibody fragments capable of binding to amyloid or portions thereof, including, but not limited to Fab (e.g., by papain digestion), Fab' (e.g., by pepsin digestion and partial reduction) and F(ab')₂ (e.g., by pepsin digestion), facb (e.g., by plasmin digestion), pFc' (e.g., by pepsin or plasmin digestion), Fd (e.g., by pepsin digestion, partial reduction and reaggregation), Fv or scFv (e.g., by molecular biology techniques) fragments, are encompassed by the invention (see, e.g., Colligan, Immunology, supra).

Such fragments can be produced by enzymatic cleavage, synthetic or recombinant techniques, as known in the art and/or as described herein. Antibodies can also be produced in a variety of truncated forms using antibody genes in which one or more stop codons have been introduced upstream of the natural stop site. For example, a combination gene encoding a F(ab')₂ heavy chain portion can be designed to include DNA sequences encoding the CH₁ domain and/or hinge region of the heavy chain. The various portions of antibodies can be joined together chemically by conventional techniques, or can be prepared as a contiguous protein using genetic engineering techniques.

As used herein, the term "human antibody" refers to an antibody in which substantially every part of the protein (e.g., CDR, framework, C_L, C_H domains (e.g., C_H1, C_H2, C_H3), hinge, (V_L, V_H)) is substantially non-immunogenic in humans, with only minor sequence changes or variations. Similarly, antibodies designated primate (monkey, baboon, chimpanzee, etc.), rodent (mouse, rat, rabbit, guinea pig, hamster, and the like) and other mammals designate such species, sub-genus, genus, sub-family, family specific antibodies. Further, chimeric antibodies of the invention can include any combination of the above. Such changes or variations optionally and preferably retain or reduce the immunogenicity in humans or other species relative to non-modified antibodies. Thus, a human antibody is distinct from a chimeric or

humanized antibody. It is pointed out that a human antibody can be produced by a non-human animal or prokaryotic or eukaryotic cell that is capable of expressing functionally rearranged human immunoglobulin (e.g., heavy chain and/or light chain) genes. Further, when a human antibody is a single chain antibody, it can comprise a linker peptide that is not found in native human antibodies. For example, an Fv can comprise a linker peptide, such as two to about eight glycine or other amino acid residues, which connects the variable region of the heavy chain and the variable region of the light chain. Such linker peptides are considered to be of human origin.

Bispecific, heterospecific, heteroconjugate or similar antibodies can also be used that are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for at least one amyloid protein, the other one is for any other antigen. Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, Nature 305:537 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed, e.g., in WO 93/08829, US Patent Nos, 6210668, 6193967, 6132992, 6106833, 6060285, 6037453, 6010902, 5989530, 5959084, 5959083, 5932448, 5833985, 5821333, 5807706, 5643759, 5601819, 5582996, 5496549, 4676980, WO 91/00360, WO 92/00373, EP 03089, Traunecker et al., EMBO J. 10:3655 (1991), Suresh et al., Methods in Enzymology 121:210 (1986), each entirely incorporated herein by reference.

Anti-amyloid antibodies (also termed amyloid antibodies) useful in the methods and compositions of the present invention can optionally be characterized by high affinity binding to amyloid and optionally and preferably having low toxicity. In particular, an antibody, specified fragment or variant of the invention, where the individual components, such as the variable region, constant region and framework, individually and/or collectively, optionally and preferably possess low immunogenicity, is useful in the present invention. The antibodies that can be used in the invention are optionally characterized by their ability to treat patients for extended periods with measurable alleviation of symptoms and low and/or acceptable toxicity. Low or acceptable immunogenicity and/or high affinity, as well as other suitable properties, can contribute to the therapeutic results achieved. "Low immunogenicity" is

defined herein as raising significant HAHA, HACA or HAMA responses in less than about 75%, or preferably less than about 50% of the patients treated and/or raising low titres in the patient treated (less than about 300, preferably less than about 100 measured with a double antigen enzyme immunoassay) (Elliott *et al.*, *Lancet* 344:1125-1127 (1994), entirely
5 incorporated herein by reference).

Utility

The isolated nucleic acids of the present invention can be used for production of at least one anti-amyloid antibody or specified variant thereof, which can be used to measure or effect in an cell, tissue, organ or animal (including mammals and humans), to diagnose, monitor,
10 modulate, treat, alleviate, help prevent the incidence of, or reduce the symptoms of, at least one amyloid condition, selected from, but not limited to, at least one of an immune disorder or disease, a cardiovascular disorder or disease, an infectious, malignant, and/or neurologic disorder or disease, or other known or specified amyloid related condition.

Such a method can comprise administering an effective amount of a composition or a
15 pharmaceutical composition comprising at least one anti-amyloid antibody to a cell, tissue, organ, animal or patient in need of such modulation, treatment, alleviation, prevention, or reduction in symptoms, effects or mechanisms. The effective amount can comprise an amount of about 0.001 to 500 mg/kg per single (e.g., bolus), multiple or continuous administration, or to achieve a serum concentration of 0.01-5000 µg/ml serum concentration per single, multiple,
20 or continuous administration, or any effective range or value therein, as done and determined using known methods, as described herein or known in the relevant arts.

Citations

All publications or patents cited herein are entirely incorporated herein by reference as they show the state of the art at the time of the present invention and/or to provide description
25 and enablement of the present invention. Publications refer to any scientific or patent publications, or any other information available in any media format, including all recorded, electronic or printed formats. The following references are entirely incorporated herein by reference: Ausubel, et al., ed., *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., NY, NY (1987-2001); Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, 2nd
30 Edition, Cold Spring Harbor, NY (1989); Harlow and Lane, *antibodies, a Laboratory Manual*, Cold Spring Harbor, NY (1989); Colligan, et al., eds., *Current Protocols in Immunology*, John Wiley & Sons, Inc., NY (1994-2001); Colligan et al., *Current Protocols in Protein Science*, John Wiley & Sons, NY, NY, (1997-2001).

Antibodies of the Present Invention

At least one anti-amyloid antibody of the present invention can be optionally produced by a cell line, a mixed cell line, an immortalized cell or clonal population of immortalized cells, as well known in the art. See, e.g., Ausubel, et al., ed., Current Protocols in Molecular Biology, John Wiley & Sons, Inc., NY, NY (1987-2001); Sambrook, et al., Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor, NY (1989); Harlow and Lane, antibodies, a Laboratory Manual, Cold Spring Harbor, NY (1989); Colligan, et al., eds., Current Protocols in Immunology, John Wiley & Sons, Inc., NY (1994-2001); Colligan et al., Current Protocols in Protein Science, John Wiley & Sons, NY, NY, (1997-2001), each entirely incorporated herein by reference.

Human antibodies that are specific for human amyloid proteins or fragments thereof can be raised against an appropriate immunogenic antigen, such as isolated and/or amyloid protein or a portion thereof (including synthetic molecules, such as synthetic peptides), e.g., but not limited to at least one of amino acid 1-7, 1-40, 31-42 and 36-40 of SEQ ID NO:50.

Other specific or general mammalian antibodies can be similarly raised. Preparation of immunogenic antigens, and monoclonal antibody production can be performed using any suitable technique.

In one approach, a hybridoma is produced by fusing a suitable immortal cell line (e.g., a myeloma cell line such as, but not limited to, Sp2/0, Sp2/0-AG14, NSO, NS1, NS2, AE-1, L.5, >243, P3X63Ag8.653, Sp2 SA3, Sp2 MAI, Sp2 SS1, Sp2 SA5, U937, MLA 144, ACT IV, MOLT4, DA-1, JURKAT, WEHI, K-562, COS, RAJI, NIH 3T3, HL-60, MLA 144, NAMAIWA, NEURO 2A, or the like, or heteromyelomas, fusion products thereof, or any cell or fusion cell derived therefrom, or any other suitable cell line as known in the art. See, e.g., www.atcc.org, www.lifetech.com., and the like, with antibody producing cells, such as, but not limited to, isolated or cloned spleen, peripheral blood, lymph, tonsil, or other immune or B cell containing cells, or any other cells expressing heavy or light chain constant or variable or framework or CDR sequences, either as endogenous or heterologous nucleic acid, as recombinant or endogenous, viral, bacterial, algal, prokaryotic, amphibian, insect, reptilian, fish, mammalian, rodent, equine, ovine, goat, sheep, primate, eukaryotic, genomic DNA, cDNA, rDNA, mitochondrial DNA or RNA, chloroplast DNA or RNA, hnRNA, mRNA, tRNA, single, double or triple stranded, hybridized, and the like or any combination thereof. See, e.g., Ausubel, supra, and Colligan, Immunology, supra, chapter 2, entirely incorporated herein by reference.

Antibody producing cells can also be obtained from the peripheral blood or, preferably the spleen or lymph nodes, of humans or other suitable animals that have been immunized with

the antigen of interest. Any other suitable host cell can also be used for expressing heterologous or endogenous nucleic acid encoding an antibody, specified fragment or variant thereof, of the present invention. The fused cells (hybridomas) or recombinant cells can be isolated using selective culture conditions or other suitable known methods, and cloned by
5 limiting dilution or cell sorting, or other known methods. Cells which produce antibodies with the desired specificity can be selected by a suitable assay (e.g., ELISA).

Other suitable methods of producing or isolating antibodies of the requisite specificity can be used, including, but not limited to, methods that select recombinant antibody from a peptide or protein library (e.g., but not limited to, a bacteriophage, ribosome, oligonucleotide,
10 RNA, cDNA, or the like, display library; e.g., as available from Cambridge antibody Technologies, Cambridgeshire, UK; MorphoSys, Martinsreid/Planegg, DE; Biovation, Aberdeen, Scotland, UK; BioInvent, Lund, Sweden; Dyax Corp., Enzon, Affymax/Biosite; Xoma, Berkeley, CA; Ixsys. See, e.g., EP 368,684, PCT/GB91/01134; PCT/GB92/01755; PCT/GB92/002240; PCT/GB92/00883; PCT/GB93/00605; US 08/350260(5/12/94);
15 PCT/GB94/01422; PCT/GB94/02662; PCT/GB97/01835; (CAT/MRC); WO90/14443; WO90/14424; WO90/14430; PCT/US94/1234; WO92/18619; WO96/07754; (Scripps); WO96/13583, WO97/08320 (MorphoSys); WO95/16027 (BioInvent); WO88/06630; WO90/3809 (Dyax); US 4,704,692 (Enzon); PCT/US91/02989 (Affymax); WO89/06283; EP 371 998; EP 550 400; (Xoma); EP 229 046; PCT/US91/07149 (Ixsys); or stochastically
20 generated peptides or proteins - US 5723323, 5763192, 5814476, 5817483, 5824514, 5976862, WO 86/05803, EP 590 689 (Ixsys, now Applied Molecular Evolution (AME), each entirely incorporated herein by reference) or that rely upon immunization of transgenic animals (e.g., SCID mice, Nguyen et al., Microbiol. Immunol. 41:901-907 (1997); Sandhu et al., Crit. Rev. Biotechnol. 16:95-118 (1996); Eren et al., Immunol. 93:154-161 (1998), each entirely
25 incorporated by reference as well as related patents and applications) that are capable of producing a repertoire of human antibodies, as known in the art and/or as described herein. Such techniques, include, but are not limited to, ribosome display (Hanes et al., Proc. Natl. Acad. Sci. USA, 94:4937-4942 (May 1997); Hanes et al., Proc. Natl. Acad. Sci. USA, 95:14130-14135 (Nov. 1998)); single cell antibody producing technologies (e.g., selected
30 lymphocyte antibody method ("SLAM") (US pat. No. 5,627,052, Wen et al., J. Immunol. 17:887-892 (1987); Babcook et al., Proc. Natl. Acad. Sci. USA 93:7843-7848 (1996)); gel microdroplet and flow cytometry (Powell et al., Biotechnol. 8:333-337 (1990); One Cell Systems, Cambridge, MA; Gray et al., J. Imm. Meth. 182:155-163 (1995); Kenny et al., Bio/Technol. 13:787-790 (1995)); B-cell selection (Steenbakketers et al., Molec. Biol. Reports 19:125-134 (1994); Jonak et al., Progress Biotech, Vol. 5, In Vitro Immunization in Hybridoma
35

Technology, Borrebaeck, ed., Elsevier Science Publishers B.V., Amsterdam, Netherlands (1988)).

Methods for engineering or humanizing non-human or human antibodies can also be used and are well known in the art. Generally, a humanized or engineered antibody has one or
5 more amino acid residues from a source which is non-human, e.g., but not limited to mouse, rat, rabbit, non-human primate or other mammal. These human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable, constant or other domain of a known human sequence.

Methods for engineering or humanizing non-human or human antibodies can also be
10 used and are well known in the art. Generally, a humanized or engineered antibody has one or more amino acid residues from a source which is non-human, e.g., but not limited to mouse, rat, rabbit, non-human primate or other mammal. These human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable, constant or other domain of a known human sequence.

15 By "humanized antibody" is meant an antibody that is composed partially or fully of amino acid sequences derived from a human antibody germline by altering the sequence of an antibody having non-human complementarity determining regions (CDR). The simplest such alteration may consist simply of substituting the constant region of a human antibody for the murine constant region, thus resulting in a human/murine chimera which may have sufficiently
20 low immunogenicity to be acceptable for pharmaceutical use.

Preferably, however, the variable region of the antibody and even the CDR is also humanized by techniques that are by now well known in the art. The framework regions of the variable regions are substituted by the corresponding human framework regions leaving the non-human CDR substantially intact, or even replacing the CDR with sequences derived from
25 a human genome. Fully human antibodies are produced in genetically modified mice whose immune systems have been altered to correspond to human immune systems. As mentioned above, it is sufficient for use in the methods of the invention, to employ an immunologically specific fragment of the antibody, including fragments representing single chain forms.

A humanized antibody again refers to an antibody comprising a human framework, at
30 least one CDR from a non-human antibody, and in which any constant region present is substantially identical to a human immunoglobulin constant region, i.e., at least about 85-90%, preferably at least 95% identical. Hence, all parts of a humanized antibody, except possibly the CDRs, are substantially identical to corresponding parts of one or more native human immunoglobulin sequences. For example, a humanized immunoglobulin would typically not

encompass a chimeric mouse variable region/human constant region antibody.

Humanized antibodies have at least three potential advantages over non-human and chimeric antibodies for use in human therapy:

1) Because the effector portion is human, it may interact better with the other parts of the human immune system (e.g., destroy the target cells more efficiently by complement-dependent cytotoxicity (CDC) or antibody-dependent cellular cytotoxicity (ADCC)).

2) The human immune system should not recognize the framework or C region of the humanized antibody as foreign, and therefore the antibody response against such an injected antibody should be less than against a totally foreign non-human antibody or a partially foreign chimeric antibody.

3) Injected non-human antibodies have been reported to have a half-life in the human circulation much shorter than the half-life of human antibodies. Injected humanized antibodies will have a half-life essentially identical to naturally occurring human antibodies, allowing smaller and less frequent doses to be given.

Known human Ig sequences are disclosed, e.g.,
www.ncbi.nlm.nih.gov/entrez/query.fcgi; www.atcc.org/phage/hdb.html; www.sciquest.com/;
www.abcam.com/; www.antibodyresource.com/onlinecomp.html;
www.public.iastate.edu/~pedro/research_tools.html; www.mgen.uni-heidelberg.de/SD/IT/IT.html; www.whfreeman.com/immunology/CH05/kuby05.htm;
www.library.thinkquest.org/12429/Immune/Antibody.html;
www.hhmi.org/grants/lectures/1996/vlab/; www.path.cam.ac.uk/~mrc7/mikeimages.html;
www.antibodyresource.com/;
mcb.harvard.edu/BioLinks/Immunology.html; www.immunologylink.com/;
pathbox.wustl.edu/~hcenter/index.html; www.biotech.ufl.edu/~hcl/;
www.pebio.com/pa/340913/340913.html; www.nal.usda.gov/awic/pubs/antibody/;
www.m.ehime-u.ac.jp/~yasuhito/Elisa.html; www.biodesign.com/table.asp;
www.icnet.uk/axp/facs/davies/links.html; www.biotech.ufl.edu/~fccl/protocol.html; www.isac-net.org/sites_geo.html; aximtl1.imt.uni-marburg.de/~rek/AEPStart.html;
baserv.uci.kun.nl/~jraats/links1.html; www.recab.uni-hd.de/immuno.bme.nwu.edu/; www.mrc-cpe.cam.ac.uk/imt-doc/public/INTRO.html; www.ibt.unam.mx/vir/V_mice.html;
imgt.cnusc.fr:8104/; www.biochem.ucl.ac.uk/~martin/abs/index.html; antibody.bath.ac.uk/;
abgen.cvm.tamu.edu/lab/wwwabgen.html;
www.unizh.ch/~honegger/AHOseminar/Slide01.html; www.cryst.bbk.ac.uk/~ubcg07s/;
www.nimr.mrc.ac.uk/CC/caewg/caewg.htm;
CEN5021 NP

www.path.cam.ac.uk/~mrc7/humanisation/TAHHP.html;
www.ibt.unam.mx/vir/structure/stat_aim.html; www.biosci.missouri.edu/smithgp/index.html;
www.cryst.bioc.cam.ac.uk/~fmolina/Web-pages/Pept/spottech.html;
www.jerini.de/fr_products.htm; www.patents.ibm.com/ibm.html. Kabat et al., Sequences of
5 Proteins of Immunological Interest, U.S. Dept. Health (1983), each entirely incorporated herein
by reference.

Such imported sequences can be used to reduce immunogenicity or reduce, enhance or
modify binding, affinity, on-rate, off-rate, avidity, specificity, half-life, or any other suitable
characteristic, as known in the art. Generally part or all of the non-human or human CDR
10 sequences are maintained while the non-human sequences of the variable and constant regions
are replaced with human or other amino acids. Antibodies can also optionally be humanized
with retention of high affinity for the antigen and other favorable biological properties. To
achieve this goal, humanized antibodies can be optionally prepared by a process of analysis of
the parental sequences and various conceptual humanized products using three-dimensional
15 models of the parental and humanized sequences. Three-dimensional immunoglobulin models
are commonly available and are familiar to those skilled in the art. Computer programs are
available which illustrate and display probable three-dimensional conformational structures of
selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of
the likely role of the residues in the functioning of the candidate immunoglobulin sequence,
20 i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind
its antigen. In this way, FR residues can be selected and combined from the consensus and
import sequences so that the desired antibody characteristic, such as increased affinity for the
target antigen(s), is achieved. In general, the CDR residues are directly and most substantially
involved in influencing antigen binding. Humanization or engineering of antibodies of the
25 present invention can be performed using any known method, such as but not limited to those
described in, Winter (Jones et al., Nature 321:522 (1986); Riechmann et al., Nature 332:323
(1988); Verhoeyen et al., Science 239:1534 (1988)), Sims et al., J. Immunol. 151: 2296 (1993);
Chothia and Lesk, J. Mol. Biol. 196:901 (1987), Carter et al., Proc. Natl. Acad. Sci. U.S.A.
89:4285 (1992); Presta et al., J. Immunol. 151:2623 (1993), US patent Nos: 5723323, 5976862,
30 5824514, 5817483, 5814476, 5763192, 5723323, 5,766886, 5714352, 6204023, 6180370,
5693762, 5530101, 5585089, 5225539; 4816567, PCT/: US98/16280, US96/18978,
US91/09630, US91/05939, US94/01234, GB89/01334, GB91/01134, GB92/01755;
WO90/14443, WO90/14424, WO90/14430, EP 229246, each entirely incorporated herein by
reference, included references cited therein.

35 The anti-amyloid antibody can also be optionally generated by immunization of a

transgenic animal (e.g., mouse, rat, hamster, non-human primate, and the like) capable of producing a repertoire of human antibodies, as described herein and/or as known in the art. Cells that produce a human anti-amyloid antibody can be isolated from such animals and immortalized using suitable methods, such as the methods described herein.

5 Transgenic mice that can produce a repertoire of human antibodies that bind to human antigens can be produced by known methods (e.g., but not limited to, U.S. Pat. Nos: 5,770,428, 5,569,825, 5,545,806, 5,625,126, 5,625,825, 5,633,425, 5,661,016 and 5,789,650 issued to Lonberg *et al.*; Jakobovits *et al.* WO 98/50433, Jakobovits *et al.* WO 98/24893, Lonberg *et al.* WO 98/24884, Lonberg *et al.* WO 97/13852, Lonberg *et al.* WO 94/25585, Kucherlapate *et al.* 10 WO 96/34096, Kucherlapate *et al.* EP 0463 151 B1, Kucherlapate *et al.* EP 0710 719 A1, Surani *et al.* US. Pat. No. 5,545,807, Bruggemann *et al.* WO 90/04036, Bruggemann *et al.* EP 0438 474 B1, Lonberg *et al.* EP 0814 259 A2, Lonberg *et al.* GB 2 272 440 A, Lonberg *et al.* *Nature* 368:856-859 (1994), Taylor *et al.*, *Int. Immunol.* 6(4):579-591 (1994), Green *et al.*, *Nature Genetics* 7:13-21 (1994), Mendez *et al.*, *Nature Genetics* 15:146-156 (1997), Taylor *et al.*, 15 *Nucleic Acids Research* 20(23):6287-6295 (1992), Tuaillon *et al.*, *Proc Natl Acad Sci USA* 90(8):3720-3724 (1993), Lonberg *et al.*, *Int Rev Immunol* 13(1):65-93 (1995) and Fishwald *et al.*, *Nat Biotechnol* 14(7):845-851 (1996), which are each entirely incorporated herein by reference). Generally, these mice comprise at least one transgene comprising DNA from at least one human immunoglobulin locus that is functionally rearranged, or which can undergo 20 functional rearrangement. The endogenous immunoglobulin loci in such mice can be disrupted or deleted to eliminate the capacity of the animal to produce antibodies encoded by endogenous genes.

Screening antibodies for specific binding to similar proteins or fragments can be conveniently achieved using peptide display libraries. This method involves the screening of 25 large collections of peptides for individual members having the desired function or structure. Antibody screening of peptide display libraries is well known in the art. The displayed peptide sequences can be from 3 to 5000 or more amino acids in length, frequently from 5-100 amino acids long, and often from about 8 to 25 amino acids long. In addition to direct chemical synthetic methods for generating peptide libraries, several recombinant DNA methods have been 30 described. One type involves the display of a peptide sequence on the surface of a bacteriophage or cell. Each bacteriophage or cell contains the nucleotide sequence encoding the particular displayed peptide sequence. Such methods are described in PCT Patent Publication Nos. 91/17271, 91/18980, 91/19818, and 93/08278. Other systems for generating libraries of peptides have aspects of both in vitro chemical synthesis and recombinant methods. See, PCT Patent 35 Publication Nos. 92/05258, 92/14843, and 96/19256. See also, U.S. Patent Nos. 5,658,754; and

5,643,768. Peptide display libraries, vector, and screening kits are commercially available from such suppliers as Invitrogen (Carlsbad, CA), and Cambridge antibody Technologies (Cambridgeshire, UK). See, e.g., U.S. Pat. Nos. 4704692, 4939666, 4946778, 5260203, 5455030, 5518889, 5534621, 5656730, 5763733, 5767260, 5856456, assigned to Enzon; 5223409, 5403484, 5571698, 5837500, assigned to Dyax, 5427908, 5580717, assigned to Affymax; 5885793, assigned to Cambridge antibody Technologies; 5750373, assigned to Genentech, 5618920, 5595898, 5576195, 5698435, 5693493, 5698417, assigned to Xoma, Colligan, *supra*; Ausubel, *supra*; or Sambrook, *supra*, each of the above patents and publications entirely incorporated herein by reference.

10 Antibodies of the present invention can also be prepared using at least one anti-amyloid antibody encoding nucleic acid to provide transgenic animals or mammals, such as goats, cows, horses, sheep, and the like, that produce such antibodies in their milk. Such animals can be provided using known methods. See, e.g., but not limited to, US patent nos. 5,827,690; 5,849,992; 4,873,316; 5,849,992; 5,994,616; 5,565,362; 5,304,489, and the like, 15 each of which is entirely incorporated herein by reference.

Antibodies of the present invention can additionally be prepared using at least one anti-amyloid antibody encoding nucleic acid to provide transgenic plants and cultured plant cells (e.g., but not limited to tobacco and maize) that produce such antibodies, specified portions or variants in the plant parts or in cells cultured therefrom. As a non-limiting 20 example, transgenic tobacco leaves expressing recombinant proteins have been successfully used to provide large amounts of recombinant proteins, e.g., using an inducible promoter. See, e.g., Cramer et al., Curr. Top. Microbol. Immunol. 240:95-118 (1999) and references cited therein. Also, transgenic maize have been used to express mammalian proteins at commercial production levels, with biological activities equivalent to those produced in other recombinant 25 systems or purified from natural sources. See, e.g., Hood et al., Adv. Exp. Med. Biol. 464:127-147 (1999) and references cited therein. antibodies have also been produced in large amounts from transgenic plant seeds including antibody fragments, such as single chain antibodies (scFv's), including tobacco seeds and potato tubers. See, e.g., Conrad et al., Plant Mol. Biol. 38:101-109 (1998) and reference cited therein. Thus, antibodies of the present invention can 30 also be produced using transgenic plants, according to know methods. See also, e.g., Fischer et al., Biotechnol. Appl. Biochem. 30:99-108 (Oct., 1999), Ma et al., Trends Biotechnol. 13:522-7 (1995); Ma et al., Plant Physiol. 109:341-6 (1995); Whitelam et al., Biochem. Soc. Trans. 22:940-944 (1994); and references cited therein. See, also generally for plant expression of antibodies, but not limited to. Each of the above references is entirely incorporated herein by 35 reference.

The antibodies of the invention can bind human amyloid with a wide range of affinities (K_D). In a preferred embodiment, at least one human mAb of the present invention can optionally bind human amyloid with high affinity. For example, a human mAb can bind human amyloid with a K_D equal to or less than about 10^{-7} M, such as but not limited to, 0.1-9.9
5 (or any range or value therein) $\times 10^{-7}$, 10^{-8} , 10^{-9} , 10^{-10} , 10^{-11} , 10^{-12} , 10^{-13} or any range or value therein.

The affinity or avidity of an antibody for an antigen can be determined experimentally using any suitable method. (See, for example, Berzofsky, *et al.*, "Antibody-Antigen Interactions," In *Fundamental Immunology*, Paul, W. E., Ed., Raven Press: New York, NY
10 (1984); Kubly, Janis *Immunology*, W. H. Freeman and Company: New York, NY (1992); and methods described herein). The measured affinity of a particular antibody-antigen interaction can vary if measured under different conditions (e.g., salt concentration, pH). Thus, measurements of affinity and other antigen-binding parameters (e.g., K_D , K_a , K_d) are preferably made with standardized solutions of antibody and antigen, and a standardized buffer, such as
15 the buffer described herein.

Nucleic Acid Molecules

Using the information provided herein, such as the nucleotide sequences encoding at least 70-100% of the contiguous amino acids of at least one of SEQ ID NOS:42-49, 53-60, 63-70, 73-80, specified fragments, variants or consensus sequences thereof, or a deposited vector
20 comprising at least one of these sequences, a nucleic acid molecule of the present invention encoding at least one anti-amyloid antibody can be obtained using methods described herein or as known in the art.

Nucleic acid molecules of the present invention can be in the form of RNA, such as mRNA, hnRNA, tRNA or any other form, or in the form of DNA, including, but not limited to,
25 cDNA and genomic DNA obtained by cloning or produced synthetically, or any combinations thereof. The DNA can be triple-stranded, double-stranded or single-stranded, or any combination thereof. Any portion of at least one strand of the DNA or RNA can be the coding strand, also known as the sense strand, or it can be the non-coding strand, also referred to as the anti-sense strand.

Isolated nucleic acid molecules of the present invention can include nucleic acid molecules comprising an open reading frame (ORF), optionally with one or more introns, e.g., but not limited to, at least one specified portion of at least one CDR, as CDR1, CDR2 and/or CDR3 of at least one heavy chain (e.g., SEQ ID NOS:42-44, 53-55, 63-65, 73-75) or light chain (e.g., SEQ ID NOS:45-47, 56-58, 66-68, 76-78); nucleic acid molecules comprising the
35 coding sequence for an anti-amyloid antibody or variable region (e.g., SEQ ID NOS:48, 49, 59,

60, 69, 70, 79 and 80), such as but not limited to SEQ ID NOS:51, 52, 61, 62, 71, 72, 81 and 82; and nucleic acid molecules which comprise a nucleotide sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode at least one anti-amyloid antibody as described herein and/or as known in the art. Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate such degenerate nucleic acid variants that code for specific anti-amyloid antibodies of the present invention. See, e.g., Ausubel, et al., *supra*, and such nucleic acid variants are included in the present invention.

As indicated herein, nucleic acid molecules of the present invention which comprise a nucleic acid encoding an anti-amyloid antibody can include, but are not limited to, those encoding the amino acid sequence of an antibody fragment, by itself; the coding sequence for the entire antibody or a portion thereof; the coding sequence for an antibody, fragment or portion, as well as additional sequences, such as the coding sequence of at least one signal leader or fusion peptide, with or without the aforementioned additional coding sequences, such as at least one intron, together with additional, non-coding sequences, including but not limited to, non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription, mRNA processing, including splicing and polyadenylation signals (for example, ribosome binding and stability of mRNA); an additional coding sequence that codes for additional amino acids, such as those that provide additional functionalities. Thus, the sequence encoding an antibody can be fused to a marker sequence, such as a sequence encoding a peptide that facilitates purification of the fused antibody comprising an antibody fragment or portion.

Polynucleotides Which Selectively Hybridize to a Polynucleotide as Described Herein

The present invention provides isolated nucleic acids that hybridize under selective hybridization conditions to a polynucleotide disclosed herein. Thus, the polynucleotides of this embodiment can be used for isolating, detecting, and/or quantifying nucleic acids comprising such polynucleotides. For example, polynucleotides of the present invention can be used to identify, isolate, or amplify partial or full-length clones in a deposited library. In some embodiments, the polynucleotides are genomic or cDNA sequences isolated, or otherwise complementary to, a cDNA from a human or mammalian nucleic acid library.

Preferably, the cDNA library comprises at least 80% full-length sequences, preferably at least 85% or 90% full-length sequences, and more preferably at least 95% full-length sequences. The cDNA libraries can be normalized to increase the representation of rare sequences. Low or moderate stringency hybridization conditions are typically, but not exclusively, employed with sequences having a reduced sequence identity relative to complementary sequences. Moderate

and high stringency conditions can optionally be employed for sequences of greater identity. Low stringency conditions allow selective hybridization of sequences having about 70% sequence identity and can be employed to identify orthologous or paralogous sequences.

Optionally, polynucleotides of this invention will encode at least a portion of an antibody
5 encoded by the polynucleotides described herein. The polynucleotides of this invention embrace nucleic acid sequences that can be employed for selective hybridization to a polynucleotide encoding an antibody of the present invention. See, e.g., Ausubel, *supra*; Colligan, *supra*, each entirely incorporated herein by reference.

Construction of Nucleic Acids

10 The isolated nucleic acids of the present invention can be made using (a) recombinant methods, (b) synthetic techniques, (c) purification techniques, or combinations thereof, as well-known in the art.

The nucleic acids can conveniently comprise sequences in addition to a polynucleotide of the present invention. For example, a multi-cloning site comprising one or more endonuclease
15 restriction sites can be inserted into the nucleic acid to aid in isolation of the polynucleotide. Also, translatable sequences can be inserted to aid in the isolation of the translated polynucleotide of the present invention. For example, a hexa-histidine marker sequence provides a convenient means to purify the proteins of the present invention. The nucleic acid of the present invention - excluding the coding sequence - is optionally a vector, adapter, or linker for cloning and/or
20 expression of a polynucleotide of the present invention.

Additional sequences can be added to such cloning and/or expression sequences to optimize their function in cloning and/or expression, to aid in isolation of the polynucleotide, or to improve the introduction of the polynucleotide into a cell. Use of cloning vectors, expression vectors, adapters, and linkers is well known in the art. (See, e.g., Ausubel, *supra*; or Sambrook,
25 *supra*)

Recombinant Methods for Constructing Nucleic Acids

The isolated nucleic acid compositions of this invention, such as RNA, cDNA, genomic DNA, or any combination thereof, can be obtained from biological sources using any number of cloning methodologies known to those of skill in the art. In some embodiments, oligonucleotide
30 probes that selectively hybridize, under stringent conditions, to the polynucleotides of the present invention are used to identify the desired sequence in a cDNA or genomic DNA library. The isolation of RNA, and construction of cDNA and genomic libraries, is well known to those of ordinary skill in the art. (See, e.g., Ausubel, *supra*; or Sambrook, *supra*)

Nucleic Acid Screening and Isolation Methods

A cDNA or genomic library can be screened using a probe based upon the sequence of a polynucleotide of the present invention, such as those disclosed herein. Probes can be used to hybridize with genomic DNA or cDNA sequences to isolate homologous genes in the same or different organisms. Those of skill in the art will appreciate that various degrees of stringency of hybridization can be employed in the assay; and either the hybridization or the wash medium can be stringent. As the conditions for hybridization become more stringent, there must be a greater degree of complementarity between the probe and the target for duplex formation to occur. The degree of stringency can be controlled by one or more of temperature, ionic strength, pH and the presence of a partially denaturing solvent such as formamide. For example, the stringency of hybridization is conveniently varied by changing the polarity of the reactant solution through, for example, manipulation of the concentration of formamide within the range of 0% to 50%. The degree of complementarity (sequence identity) required for detectable binding will vary in accordance with the stringency of the hybridization medium and/or wash medium. The degree of complementarity will optimally be 100%, or 70-100%, or any range or value therein. However, it should be understood that minor sequence variations in the probes and primers can be compensated for by reducing the stringency of the hybridization and/or wash medium.

Methods of amplification of RNA or DNA are well known in the art and can be used according to the present invention without undue experimentation, based on the teaching and guidance presented herein.

Known methods of DNA or RNA amplification include, but are not limited to, polymerase chain reaction (PCR) and related amplification processes (see, e.g., U.S. Patent Nos. 4,683,195, 4,683,202, 4,800,159, 4,965,188, to Mullis, et al.; 4,795,699 and 4,921,794 to Tabor, et al; 5,142,033 to Innis; 5,122,464 to Wilson, et al.; 5,091,310 to Innis; 5,066,584 to Gyllenstein, et al; 4,889,818 to Gelfand, et al; 4,994,370 to Silver, et al; 4,766,067 to Biswas; 4,656,134 to Ringold) and RNA mediated amplification that uses anti-sense RNA to the target sequence as a template for double-stranded DNA synthesis (U.S. Patent No. 5,130,238 to Malek, et al, with the tradename NASBA), the entire contents of which references are incorporated herein by reference. (See, e.g., Ausubel, *supra*; or Sambrook, *supra*.)

For instance, polymerase chain reaction (PCR) technology can be used to amplify the sequences of polynucleotides of the present invention and related genes directly from genomic DNA or cDNA libraries. PCR and other in vitro amplification methods can also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes. Examples of techniques sufficient to direct persons of skill through in vitro amplification methods are found in Berger, *supra*, Sambrook, *supra*, and

Ausubel, supra, as well as Mullis, et al., U.S. Patent No. 4,683,202 (1987); and Innis, et al., PCR Protocols A Guide to Methods and Applications, Eds., Academic Press Inc., San Diego, CA (1990). Commercially available kits for genomic PCR amplification are known in the art. See, e.g., Advantage-GC Genomic PCR Kit (Clontech). Additionally, e.g., the T4 gene 32 protein
5 (Boehringer Mannheim) can be used to improve yield of long PCR products.

Synthetic Methods for Constructing Nucleic Acids

The isolated nucleic acids of the present invention can also be prepared by direct chemical synthesis by known methods (see, e.g., Ausubel, et al., supra). Chemical synthesis generally produces a single-stranded oligonucleotide, which can be converted into double-
10 stranded DNA by hybridization with a complementary sequence, or by polymerization with a DNA polymerase using the single strand as a template. One of skill in the art will recognize that while chemical synthesis of DNA can be limited to sequences of about 100 or more bases, longer sequences can be obtained by the ligation of shorter sequences.

Recombinant Expression Cassettes

15 The present invention further provides recombinant expression cassettes comprising a nucleic acid of the present invention. A nucleic acid sequence of the present invention, for example a cDNA or a genomic sequence encoding an antibody of the present invention, can be used to construct a recombinant expression cassette that can be introduced into at least one desired host cell. A recombinant expression cassette will typically comprise a polynucleotide of
20 the present invention operably linked to transcriptional initiation regulatory sequences that will direct the transcription of the polynucleotide in the intended host cell. Both heterologous and non-heterologous (i.e., endogenous) promoters can be employed to direct expression of the nucleic acids of the present invention.

In some embodiments, isolated nucleic acids that serve as promoter, enhancer, or other
25 elements can be introduced in the appropriate position (upstream, downstream or in intron) of a non-heterologous form of a polynucleotide of the present invention so as to up or down regulate expression of a polynucleotide of the present invention. For example, endogenous promoters can be altered *in vivo* or *in vitro* by mutation, deletion and/or substitution.

Vectors And Host Cells

30 The present invention also relates to vectors that include isolated nucleic acid molecules of the present invention, host cells that are genetically engineered with the recombinant vectors, and the production of at least one anti-amyloid antibody by recombinant techniques, as is well known in the art. See, e.g., Sambrook, et al., supra; Ausubel, et al., supra, each entirely incorporated herein by reference.

The polynucleotides can optionally be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it can be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

The DNA insert should be operatively linked to an appropriate promoter. The expression constructs will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts expressed by the constructs will preferably include a translation initiating at the beginning and a termination codon (e.g., UAA, UGA or UAG) appropriately positioned at the end of the mRNA to be translated, with UAA and UAG preferred for mammalian or eukaryotic cell expression.

Expression vectors will preferably but optionally include at least one selectable marker. Such markers include, e.g., but not limited to, methotrexate (MTX), dihydrofolate reductase (DHFR, US Pat.Nos. 4,399,216; 4,634,665; 4,656,134; 4,956,288; 5,149,636; 5,179,017, ampicillin, neomycin (G418), mycophenolic acid, or glutamine synthetase (GS, US Pat.Nos. 5,122,464; 5,770,359; 5,827,739) resistance for eukaryotic cell culture, and tetracycline or ampicillin resistance genes for culturing in *E. coli* and other bacteria or prokaryotics (the above patents are entirely incorporated hereby by reference). Appropriate culture mediums and conditions for the above-described host cells are known in the art. Suitable vectors will be readily apparent to the skilled artisan. Introduction of a vector construct into a host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other known methods. Such methods are described in the art, such as Sambrook, supra, Chapters 1-4 and 16-18; Ausubel, supra, Chapters 1, 9, 13, 15, 16.

At least one antibody of the present invention can be expressed in a modified form, such as a fusion protein, and can include not only secretion signals, but also additional heterologous functional regions. For instance, a region of additional amino acids, particularly charged amino acids, can be added to the N-terminus of an antibody to improve stability and persistence in the host cell, during purification, or during subsequent handling and storage. Also, peptide moieties can be added to an antibody of the present invention to facilitate purification. Such regions can be removed prior to final preparation of an antibody or at least one fragment thereof. Such methods are described in many standard laboratory manuals, such as Sambrook, supra, Chapters 17.29-17.42 and 18.1-18.74; Ausubel, supra, Chapters 16, 17 and 18.

Those of ordinary skill in the art are knowledgeable in the numerous expression systems available for expression of a nucleic acid encoding a protein of the present invention.

Alternatively, nucleic acids of the present invention can be expressed in a host cell by turning on (by manipulation) in a host cell that contains endogenous DNA encoding an antibody
5 of the present invention. Such methods are well known in the art, e.g., as described in US patent Nos. 5,580,734, 5,641,670, 5,733,746, and 5,733,761, entirely incorporated herein by reference.

Illustrative of cell cultures useful for the production of the antibodies, specified portions or variants thereof, are mammalian cells. Mammalian cell systems often will be in the form of monolayers of cells although mammalian cell suspensions or bioreactors can also be used. A
10 number of suitable host cell lines capable of expressing intact glycosylated proteins have been developed in the art, and include the COS-1 (e.g., ATCC CRL 1650), COS-7 (e.g., ATCC CRL-1651), HEK293, BHK21 (e.g., ATCC CRL-10), CHO (e.g., ATCC CRL 1610) and BSC-1 (e.g., ATCC CRL-26) cell lines, Cos-7 cells, CHO cells, hep G2 cells, P3X63Ag8.653, SP2/0-Ag14, 293 cells, HeLa cells and the like, which are readily available from, for example, American
15 Type Culture Collection, Manassas, Va (www.atcc.org). Host cells include cells of lymphoid origin such as myeloma and lymphoma cells. Host cells are P3X63Ag8.653 cells (ATCC Accession Number CRL-1580) and SP2/0-Ag14 cells (ATCC Accession Number CRL-1851). In a particularly preferred embodiment, the recombinant cell is a P3X63Ab8.653 or an SP2/0-Ag14 cell.

20 Expression vectors for these cells can include one or more of the following expression control sequences, such as, but not limited to an origin of replication; a promoter (e.g., late or early SV40 promoters, the CMV promoter (US Pat.Nos. 5,168,062; 5,385,839), an HSV tk promoter, a pgk (phosphoglycerate kinase) promoter, an EF-1 alpha promoter (US Pat.No. 5,266,491), at least one human immunoglobulin promoter; an enhancer, and/or processing
25 information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites (e.g., an SV40 large T Ag poly A addition site), and transcriptional terminator sequences. See, e.g., Ausubel et al., supra; Sambrook, et al., supra. Other cells useful for production of nucleic acids or proteins of the present invention are known and/or available, for instance, from the American Type Culture Collection Catalogue of Cell Lines and Hybridomas (www.atcc.org) or other
30 known or commercial sources.

When eukaryotic host cells are employed, polyadenylation or transcription terminator sequences are typically incorporated into the vector. An example of a terminator sequence is the polyadenylation sequence from the bovine growth hormone gene. Sequences for accurate splicing of the transcript can also be included. An example of a splicing sequence is the VP1

intron from SV40 (Sprague, et al., J. Virol. 45:773-781 (1983)). Additionally, gene sequences to control replication in the host cell can be incorporated into the vector, as known in the art.

Purification of an Antibody

An anti-amyloid antibody can be recovered and purified from recombinant cell
5 cultures by well-known methods including, but not limited to, protein A purification,
ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange
chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography,
affinity chromatography, hydroxylapatite chromatography and lectin chromatography. High
performance liquid chromatography ("HPLC") can also be employed for purification. See,
10 e.g., Colligan, Current Protocols in Immunology, or Current Protocols in Protein Science, John
Wiley & Sons, NY, NY, (1997-2001), e.g., Chapters 1, 4, 6, 8, 9, 10, each entirely incorporated
herein by reference.

Antibodies of the present invention include naturally purified products, products of
chemical synthetic procedures, and products produced by recombinant techniques from a
15 eukaryotic host, including, for example, yeast, higher plant, insect and mammalian cells.
Depending upon the host employed in a recombinant production procedure, the antibody of the
present invention can be glycosylated or can be non-glycosylated, with glycosylated preferred.
Such methods are described in many standard laboratory manuals, such as Sambrook, supra,
Sections 17.37-17.42; Ausubel, supra, Chapters 10, 12, 13, 16, 18 and 20, Colligan, Protein
20 Science, supra, Chapters 12-14, all entirely incorporated herein by reference.

Anti-amyloid Antibodies

The isolated antibodies of the present invention comprise an antibody amino acid
sequences disclosed herein encoded by any suitable polynucleotide, or any isolated or prepared
antibody. Preferably, the human antibody or antigen-binding fragment binds human amyloid
25 and, thereby partially or substantially neutralizes at least one biological activity of the protein.
An antibody, or specified portion or variant thereof, that partially or preferably substantially
neutralizes at least one biological activity of at least one amyloid protein or fragment can bind the
protein or fragment and thereby inhibit activitys mediated through the binding of amyloid to the
amyloid receptor or through other amyloid-dependent or mediated mechanisms. As used herein,
30 the term "neutralizing antibody" refers to an antibody that can inhibit an amyloid-dependent
activity by about 20-120%, preferably by at least about 10, 20, 30, 40, 50, 55, 60, 65, 70, 75, 80,
85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100% or more depending on the assay. The capacity of
an anti-amyloid antibody to inhibit an amyloid-dependent activity is preferably assessed by at
least one suitable amyloid protein or receptor assay, as described herein and/or as known in the

art. A human antibody of the invention can be of any class (IgG, IgA, IgM, IgE, IgD, etc.) or isotype and can comprise a kappa or lambda light chain. In one embodiment, the human antibody comprises an IgG heavy chain or defined fragment, for example, at least one of isotypes, IgG1, IgG2, IgG3 or IgG4. Antibodies of this type can be prepared by employing a transgenic mouse or
5 other transgenic non-human mammal comprising at least one human light chain (e.g., IgG, IgA and IgM (e.g., γ 1, γ 2, γ 3, γ 4) transgenes as described herein and/or as known in the art. In another embodiment, the anti-human amyloid human antibody comprises an IgG1 heavy chain and an IgG1 light chain.

At least one antibody of the invention binds at least one specified epitope specific to at
10 least one amyloid protein, subunit, fragment, portion or any combination thereof. The at least one epitope can comprise at least one antibody binding region that comprises at least one portion of the protein, which epitope is preferably comprised of at least one extracellular, soluble, hydrophilic, external or cytoplasmic portion of the protein. The at least one specified epitope can comprise any combination of at least one amino acid sequence of at least 1-3
15 amino acids to the entire specified portion of contiguous amino acids of the SEQ ID NO:50. As non-limiting examples, antibodies of the present invention showed binding of amino acids 2-7, 3-8, 33-42, and/or 34-40 of SEQ ID NO:50.

Generally, the human antibody or antigen-binding fragment of the present invention will comprise an antigen-binding region that comprises at least one human complementarity
20 determining region (CDR1, CDR2 and CDR3) or variant of at least one heavy chain variable region and at least one human complementarity determining region (CDR1, CDR2 and CDR3) or variant of at least one light chain variable region. As a non-limiting example, the antibody or antigen-binding portion or variant can comprise at least one of the heavy chain CDR3 having the amino acid sequence of SEQ ID NO:44, and/or a light chain CDR3 having the amino acid
25 sequence of SEQ ID NO:47. In a particular embodiment, the antibody or antigen-binding fragment can have an antigen-binding region that comprises at least a portion of at least one heavy chain CDR (i.e., CDR1, CDR2 and/or CDR3) having the amino acid sequence of the corresponding CDRs 1, 2 and/or 3 (e.g., SEQ ID NOS:42, 43 and/or 44; 53, 54 and/or 55; 63, 64 and/or 65; 73, 74 and/or 75). In another particular embodiment, the antibody or antigen-
30 binding portion or variant can have an antigen-binding region that comprises at least a portion of at least one light chain CDR (i.e., CDR1, CDR2 and/or CDR3) having the amino acid sequence of the corresponding CDRs 1, 2 and/or 3 (e.g., SEQ ID NOS:45, 46 and/or 47; 56, 57 and/or 58; 66, 67 and/or 68; 76, 77 and/or 78). In a preferred embodiment the three heavy chain CDRs and the three light chain CDRs of the antibody or antigen-binding fragment have
35 the amino acid sequence of the corresponding CDRs of at least one of mAb C701, C705, C706, CEN5021 NP

and C707, as described herein. Such antibodies can be prepared by chemically joining together the various portions (e.g., CDRs, framework) of the antibody using conventional techniques, by preparing and expressing a (i.e., one or more) nucleic acid molecule that encodes the antibody using conventional techniques of recombinant DNA technology or by using any other suitable method.

The anti-amyloid antibody can comprise at least one of a heavy or light chain variable region having a defined amino acid sequence. Any suitable Ig variable sequence can be used, e.g., from any subclass or any combination or fragment thereof. Such sequences are well known in the art.

As a non-limiting example, representative variable sequences include those from IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgM, and the like, e.g., HC and LC, FR1, FR2, and/or FR3 sequences from any combination of Ig subclasses, e.g., as presented in SEQ ID NOS: 48-49, 59-60, 69-70, and 79-80.

As a further non-limiting example, in a preferred embodiment, the anti-amyloid antibody comprises at least one of at least one heavy chain variable region, optionally having the amino acid sequence of SEQ ID NO:48 and/or at least one light chain variable region, optionally having the amino acid sequence of SEQ ID NO:49. In another preferred embodiment, the anti-amyloid antibody comprises at least one of at least one heavy chain variable region, optionally having the amino acid sequence of SEQ ID NO:59 and/or at least one light chain variable region, optionally having the amino acid sequence of SEQ ID NO:60. In yet another preferred embodiment, the anti-amyloid antibody comprises at least one of at least one heavy chain variable region, optionally having the amino acid sequence of SEQ ID NO:69 and/or at least one light chain variable region, optionally having the amino acid sequence of SEQ ID NO:70. In still another preferred embodiment, the anti-amyloid antibody comprises at least one of at least one heavy chain variable region, optionally having the amino acid sequence of SEQ ID NO:79 and/or at least one light chain variable region, optionally having the amino acid sequence of SEQ ID NO:80.

Antibodies that bind to human amyloid and that comprise a defined heavy or light chain variable region can be prepared using suitable methods, such as phage display (Katsube, Y., *et al.*, *Int J Mol. Med*, 1(5):863-868 (1998)) or methods that employ transgenic animals, as known in the art and/or as described herein. For example, a transgenic mouse, comprising a functionally rearranged human immunoglobulin heavy chain transgene and a transgene comprising DNA from a human immunoglobulin light chain locus that can undergo functional rearrangement, can be immunized with human amyloid or a fragment thereof to elicit the production of antibodies. If desired, the antibody producing cells can be isolated and

hybridomas or other immortalized antibody-producing cells can be prepared as described herein and/or as known in the art. Alternatively, the antibody, specified portion or variant can be expressed using the encoding nucleic acid or portion thereof in a suitable host cell.

The invention also relates to antibodies, antigen-binding fragments, immunoglobulin chains and CDRs comprising amino acids in a sequence that is substantially the same as an amino acid sequence described herein. Preferably, such antibodies or antigen-binding fragments and antibodies comprising such chains or CDRs can bind human amyloid with high affinity (e.g., K_D less than or equal to about 10^{-9} M). Amino acid sequences that are substantially the same as the sequences described herein include sequences comprising conservative amino acid substitutions, as well as amino acid deletions and/or insertions. A conservative amino acid substitution refers to the replacement of a first amino acid by a second amino acid that has chemical and/or physical properties (e.g, charge, structure, polarity, hydrophobicity/ hydrophilicity) that are similar to those of the first amino acid. Conservative substitutions include replacement of one amino acid by another within the following groups: lysine (K), arginine (R) and histidine (H); aspartate (D) and glutamate (E); asparagine (N), glutamine (Q), serine (S), threonine (T), tyrosine (Y), K, R, H, D and E; alanine (A), valine (V), leucine (L), isoleucine (I), proline (P), phenylalanine (F), tryptophan (W), methionine (M), cysteine (C) and glycine (G); F, W and Y; C, S and T.

Amino Acid Codes

The amino acids that make up anti-amyloid antibodies of the present invention are often abbreviated. The amino acid designations can be indicated by designating the amino acid by its single letter code, its three letter code, name, or three nucleotide codon(s) as is well understood in

the art (see Alberts, B., et al., Molecular Biology of The Cell, Third Ed., Garland Publishing, Inc., New York, 1994):

| SINGLE LETTER CODE | THREE LETTER CODE | NAME | THREE NUCLEOTIDE CODON(S) |
|--------------------|-------------------|---------------|------------------------------|
| A | Ala | Alanine | GCA, GCC, GCG, GCU |
| C | Cys | Cysteine | UGC, UGU |
| D | Asp | Aspartic acid | GAC, GAU |
| E | Glu | Glutamic acid | GAA, GAG |
| F | Phe | Phenylalanine | UUC, UUU |
| G | Gly | Glycine | GGA, GGC, GGG, GGU |
| H | His | Histidine | CAC, CAU |
| I | Ile | Isoleucine | AUA, AUC, AUU |
| K | Lys | Lysine | AAA, AAG |
| L | Leu | Leucine | UUA, UUG, CUA, CUC, CUG, CUU |
| M | Met | Methionine | AUG |
| N | Asn | Asparagine | AAC, AAU |
| P | Pro | Proline | CCA, CCC, CCG, CCU |
| Q | Gln | Glutamine | CAA, CAG |
| R | Arg | Arginine | AGA, AGG, CGA, CGC, CGG, CGU |
| S | Ser | Serine | AGC, AGU, UCA, UCC, UCG, UCU |
| T | Thr | Threonine | ACA, ACC, ACG, ACU |
| V | Val | Valine | GUA, GUC, GUG, GUU |
| W | Trp | Tryptophan | UGG |
| Y | Tyr | Tyrosine | UAC, UAU |

5 An anti-amyloid antibody of the present invention can include one or more amino acid substitutions, deletions or additions, either from natural mutations or human manipulation, as specified herein.

 Of course, the number of amino acid substitutions a skilled artisan would make depends on many factors, including those described above. Generally speaking, the number of
10 amino acid substitutions, insertions or deletions for any given anti-amyloid antibody, fragment or variant will not be more than 40, 30, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, such as 1-30 or any range or value therein, as specified herein.

 Amino acids in an anti-amyloid antibody of the present invention that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or
15 alanine-scanning mutagenesis (e.g., Ausubel, supra, Chapters 8, 15; Cunningham and Wells, Science 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological

activity, such as, but not limited to at least one amyloid neutralizing activity. Sites that are critical for antibody binding can also be identified by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith, et al., J. Mol. Biol. 224:899-904 (1992) and de Vos, et al., Science 255:306-312 (1992)).

5 Anti-amyloid antibodies of the present invention can include, but are not limited to, at least one portion, sequence or combination selected from 5 to all of the contiguous amino acids of at least one of SEQ ID NOS:42-47, 53-58, 63-68, or 73-78.

 An anti-amyloid antibody can further optionally comprise a polypeptide of at least one of 70-100% of the contiguous amino acids of at least one of SEQ ID NOS:48, 49, 59, 60, 69,
10 70, 79 and 80.

 In one embodiment, the amino acid sequence of an immunoglobulin chain, or portion thereof (e.g., variable region, CDR) has about 70-100% identity (e.g., 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or any range or value therein) to the amino acid sequence of the corresponding chain of at least
15 one of SEQ ID NOS:48, 49, 59, 60, 69, 70, 79 and 80. For example, the amino acid sequence of a light chain variable region can be compared with the sequence of SEQ ID NO:49, 60, 70 or 80, or the amino acid sequence of a heavy chain CDR3 can be compared with SEQ ID
NO:48, 59, 69 or 79. Preferably, 70-100% amino acid identity (i.e., 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or
20 any range or value therein) is determined using a suitable computer algorithm, as known in the art.

 Exemplary heavy chain and light chain variable regions sequences are provided in SEQ ID NOS:48, 49, 59, 60, 69, 70, 79 and 80. The antibodies of the present invention, or specified variants thereof, can comprise any number of contiguous amino acid residues from an antibody of
25 the present invention, wherein that number is selected from the group of integers consisting of from 10-100% of the number of contiguous residues in an anti-amyloid antibody. Optionally, this subsequence of contiguous amino acids is at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250 or more amino acids in length, or any range or value therein. Further, the number of such subsequences can be
30 any integer selected from the group consisting of from 1 to 20, such as at least 2, 3, 4, or 5.

 As those of skill will appreciate, the present invention includes at least one biologically active antibody of the present invention. Biologically active antibodies have a specific activity at least 20%, 30%, or 40%, and preferably at least 50%, 60%, or 70%, and most preferably at least 80%, 90%, or 95%-1000% of that of the native (non-synthetic), endogenous or related and known

antibody. Methods of assaying and quantifying measures of enzymatic activity and substrate specificity, are well known to those of skill in the art.

MODIFIED ANTIBODIES

In another aspect, the invention relates to human antibodies and antigen-binding
5 fragments, as described herein, which are modified by the covalent attachment of an organic moiety. Such modification can produce an antibody or antigen-binding fragment with improved pharmacokinetic properties (e.g., increased *in vivo* serum half-life). The organic moiety can be a linear or branched hydrophilic polymeric group, fatty acid group, or fatty acid ester group. In particular embodiments, the hydrophilic polymeric group can have a molecular
10 weight of about 800 to about 120,000 Daltons and can be a polyalkane glycol (e.g., polyethylene glycol (PEG), polypropylene glycol (PPG)), carbohydrate polymer, amino acid polymer or polyvinyl pyrrolidone, and the fatty acid or fatty acid ester group can comprise from about eight to about forty carbon atoms.

The modified antibodies and antigen-binding fragments of the invention can comprise
15 one or more organic moieties that are covalently bonded, directly or indirectly, to the antibody.

Each organic moiety that is bonded to an antibody or antigen-binding fragment of the invention can independently be a hydrophilic polymeric group, a fatty acid group or a fatty acid ester group. As used herein, the term "fatty acid" encompasses mono-carboxylic acids and di-carboxylic acids. A "hydrophilic polymeric group," as the term is used herein, refers to
20 an organic polymer that is more soluble in water than in octane. For example, polylysine is more soluble in water than in octane. Thus, an antibody modified by the covalent attachment of polylysine is encompassed by the invention. Hydrophilic polymers suitable for modifying antibodies of the invention can be linear or branched and include, for example, polyalkane glycols (e.g., PEG, monomethoxy-polyethylene glycol (mPEG), PPG and the like),
25 carbohydrates (e.g., dextran, cellulose, oligosaccharides, polysaccharides and the like), polymers of hydrophilic amino acids (e.g., polylysine, polyarginine, polyaspartate and the like), polyalkane oxides (e.g., polyethylene oxide, polypropylene oxide and the like) and polyvinyl pyrrolidone. Preferably, the hydrophilic polymer that modifies the antibody of the invention has a molecular weight of about 800 to about 150,000 Daltons as a separate
30 molecular entity. For example PEG₅₀₀₀ and PEG_{20,000}, wherein the subscript is the average molecular weight of the polymer in Daltons, can be used. The hydrophilic polymeric group can be substituted with one to about six alkyl, fatty acid or fatty acid ester groups. Hydrophilic polymers that are substituted with a fatty acid or fatty acid ester group can be prepared by employing suitable methods. For example, a polymer comprising an amine group can be
35 coupled to a carboxylate of the fatty acid or fatty acid ester, and an activated carboxylate (e.g.,

activated with N, N-carbonyl diimidazole) on a fatty acid or fatty acid ester can be coupled to a hydroxyl group on a polymer.

Fatty acids and fatty acid esters suitable for modifying antibodies of the invention can be saturated or can contain one or more units of unsaturation. Fatty acids that are suitable for
5 modifying antibodies of the invention include, for example, n-dodecanoate (C₁₂, laurate), n-tetradecanoate (C₁₄, myristate), n-octadecanoate (C₁₈, stearate), n-eicosanoate (C₂₀, arachidate), n-docosanoate (C₂₂, behenate), n-triacontanoate (C₃₀), n-tetracontanoate (C₄₀), *cis*-Δ⁹-octadecanoate (C₁₈, oleate), all *cis*-Δ^{5,8,11,14}-eicosatetraenoate (C₂₀, arachidonate), octanedioic acid, tetradecanedioic acid, octadecanedioic acid, docosanedioic acid, and the like.
10 Suitable fatty acid esters include mono-esters of dicarboxylic acids that comprise a linear or branched lower alkyl group. The lower alkyl group can comprise from one to about twelve, preferably one to about six, carbon atoms.

The modified human antibodies and antigen-binding fragments can be prepared using suitable methods, such as by reaction with one or more modifying agents. A "modifying
15 agent" as the term is used herein, refers to a suitable organic group (e.g., hydrophilic polymer, a fatty acid, a fatty acid ester) that comprises an activating group. An "activating group" is a chemical moiety or functional group that can, under appropriate conditions, react with a second chemical group thereby forming a covalent bond between the modifying agent and the second
20 chemical group. For example, amine-reactive activating groups include electrophilic groups such as tosylate, mesylate, halo (chloro, bromo, fluoro, iodo), N-hydroxysuccinimidyl esters (NHS), and the like. Activating groups that can react with thiols include, for example, maleimide, iodoacetyl, acryloyl, pyridyl disulfides, 5-thiol-2-nitrobenzoic acid thiol (TNB-thiol), and the like. An aldehyde functional group can be coupled to amine- or hydrazide-containing molecules, and an azide group can react with a trivalent phosphorous group to form
25 phosphoramidate or phosphorimide linkages. Suitable methods to introduce activating groups into molecules are known in the art (see for example, Hermanson, G. T., *Bioconjugate Techniques*, Academic Press: San Diego, CA (1996)). An activating group can be bonded directly to the organic group (e.g., hydrophilic polymer, fatty acid, fatty acid ester), or through a linker moiety, for example a divalent C₁-C₁₂ group wherein one or more carbon atoms can be
30 replaced by a heteroatom such as oxygen, nitrogen or sulfur. Suitable linker moieties include, for example, tetraethylene glycol, -(CH₂)₃-, -NH-(CH₂)₆-NH-, -(CH₂)₂-NH- and -CH₂-O-CH₂-CH₂-O-CH₂-CH₂-O-CH₂-CH₂-O-CH₂-NH-. Modifying agents that comprise a linker moiety can be produced, for example, by reacting a mono-Boc-alkyldiamine (e.g., mono-Boc-ethylenediamine, mono-Boc-diaminohexane) with a fatty acid in the presence of 1-ethyl-3-(3-
35 dimethylaminopropyl) carbodiimide (EDC) to form an amide bond between the free amine and

the fatty acid carboxylate. The Boc protecting group can be removed from the product by treatment with trifluoroacetic acid (TFA) to expose a primary amine that can be coupled to another carboxylate as described, or can be reacted with maleic anhydride and the resulting product cyclized to produce an activated maleimido derivative of the fatty acid. (See, for example, Thompson, *et al.*, WO 92/16221 the entire teachings of which are incorporated herein by reference.)

The modified antibodies of the invention can be produced by reacting a human antibody or antigen-binding fragment with a modifying agent. For example, the organic moieties can be bonded to the antibody in a non-site specific manner by employing an amine-reactive modifying agent, for example, an NHS ester of PEG. Modified human antibodies or antigen-binding fragments can also be prepared by reducing disulfide bonds (e.g., intra-chain disulfide bonds) of an antibody or antigen-binding fragment. The reduced antibody or antigen-binding fragment can then be reacted with a thiol-reactive modifying agent to produce the modified antibody of the invention. Modified human antibodies and antigen-binding fragments comprising an organic moiety that is bonded to specific sites of an antibody of the present invention can be prepared using suitable methods, such as reverse proteolysis (Fisch *et al.*, *Bioconjugate Chem.*, 3:147-153 (1992); Werlen *et al.*, *Bioconjugate Chem.*, 5:411-417 (1994); Kumaran *et al.*, *Protein Sci.* 6(10):2233-2241 (1997); Itoh *et al.*, *Bioorg. Chem.*, 24(1): 59-68 (1996); Capellas *et al.*, *Biotechnol. Bioeng.*, 56(4):456-463 (1997)), and the methods described in Hermanson, G. T., *Bioconjugate Techniques*, Academic Press: San Diego, CA (1996).

ANTI-IDIOTYPE ANTIBODIES TO ANTI-AMYLOID ANTIBODY COMPOSITIONS

In addition to monoclonal or chimeric anti-amyloid antibodies, the present invention is also directed to an anti-idiotypic (anti-Id) antibody specific for such antibodies of the invention. An anti-Id antibody is an antibody which recognizes unique determinants generally associated with the antigen-binding region of another antibody. The anti-Id can be prepared by immunizing an animal of the same species and genetic type (e.g. mouse strain) as the source of the Id antibody with the antibody or a CDR containing region thereof. The immunized animal will recognize and respond to the idiotypic determinants of the immunizing antibody and produce an anti-Id antibody. The anti-Id antibody may also be used as an "immunogen" to induce an immune response in yet another animal, producing a so-called anti-anti-Id antibody.

AMYLOID ANTIBODY COMPOSITIONS

The present invention also provides at least one anti-amyloid antibody composition comprising at least one, at least two, at least three, at least four, at least five, at least six or more anti-amyloid antibodies thereof, as described herein and/or as known in the art that are

provided in a non-naturally occurring composition, mixture or form. Such compositions comprise non-naturally occurring compositions comprising at least one or two full length, C- and/or N-terminally deleted variants, domains, fragments, or specified variants, of the anti-amyloid antibody amino acid sequence selected from the group consisting of 70-100% of the contiguous amino acids of SEQ ID NOS:42-49, 53-60, 63-70, 73-80, or specified fragments, domains or variants thereof. Preferred anti-amyloid antibody compositions include at least one or two full length, fragments, domains or variants as at least one CDR or LBP containing portions of the anti-amyloid antibody sequence of 70-100% of SEQ ID NOS:42-47, 53-58, 63-68, 73-78, or specified fragments, domains or variants thereof. Further preferred compositions comprise 40-99% of at least one of 70-100% of SEQ ID NOS:42-47, 53-58, 63-68, 73-78, or specified fragments, domains or variants thereof. Such composition percentages are by weight, volume, concentration, molarity, or molality as liquid or dry solutions, mixtures, suspension, emulsions, particles, powder, or colloids, as known in the art or as described herein.

The composition can optionally further comprise an effective amount of at least one compound or protein selected from at least one of an anti-infective drug, a cardiovascular (CV) system drug, a central nervous system (CNS) drug, an autonomic nervous system (ANS) drug, a respiratory tract drug, a gastrointestinal (GI) tract drug, a hormonal drug, a drug for fluid or electrolyte balance, a hematologic drug, an antineoplastic, an immunomodulation drug, an ophthalmic, otic or nasal drug, a topical drug, a nutritional drug, a statin, or the like. Such drugs are well known in the art, including formulations, indications, dosing and administration for each presented herein (see, e.g., Nursing 2001 Handbook of Drugs, 21st edition, Springhouse Corp., Springhouse, PA, 2001; Health Professional's Drug Guide 2001, ed., Shannon, Wilson, Stang, Prentice-Hall, Inc, Upper Saddle River, NJ; Pharmacotherapy Handbook, Wells et al., ed., Appleton & Lange, Stamford, CT, each entirely incorporated herein by reference).

The CNS drug can be at least one selected from nonnarcotic analgesics or at least one selected from antipyretics, nonsteroidal anti-inflammatory drugs, narcotic or at least one opioid analgesics, sedative-hypnotics, anticonvulsants, antidepressants, antianxiety drugs, antipsychotics, central nervous system stimulants, antiparkinsonians, miscellaneous central nervous system drugs. The ANS drug can be at least one selected from cholinergics (parasympathomimetics), anticholinergics, adrenergics (sympathomimetics), adrenergic blockers (sympatholytics), skeletal muscle relaxants, neuromuscular blockers. The at least one nonnarcotic analgesic or antipyretic can be at least one selected from acetaminophen, aspirin, choline magnesium trisalicylate, diflunisal, magnesium salicylate. The at least one nonsteroidal anti-inflammatory drug can be at least one selected from celecoxib, diclofenac

potassium, diclofenac sodium, etodolac, fenoprofen calcium, flurbiprofen, ibuprofen, indomethacin, indomethacin sodium trihydrate, ketoprofen, ketorolac tromethamine, nabumetone, naproxen, naproxen sodium, oxaprozin, piroxicam, rofecoxib, sulindac. The at least one narcotic or opioid analgesic can be at least one selected from alfentanil hydrochloride, 5 buprenorphine hydrochloride, butorphanol tartrate, codeine phosphate, codeine sulfate, fentanyl citrate, fentanyl transdermal system, fentanyl transmucosal, hydromorphone hydrochloride, meperidine hydrochloride, methadone hydrochloride, morphine hydrochloride, morphine sulfate, morphine tartrate, nalbuphine hydrochloride, oxycodone hydrochloride, oxycodone pectinate, oxymorphone hydrochloride, pentazocine hydrochloride, pentazocine 10 hydrochloride and naloxone hydrochloride, pentazocine lactate, propoxyphene hydrochloride, propoxyphene napsylate, remifentanil hydrochloride, sufentanil citrate, tramadol hydrochloride. The at least one sedative-hypnotic can be at least one selected from chloral hydrate, estazolam, flurazepam hydrochloride, pentobarbital, pentobarbital sodium, phenobarbital sodium, secobarbital sodium, temazepam, triazolam, zaleplon, zolpidem tartrate. 15 The at least one anticonvulsant can be at least one selected from acetazolamide sodium, carbamazepine, clonazepam, clorazepate dipotassium, diazepam, divalproex sodium, ethosuximide, fosphenytoin sodium, gabapentin, lamotrigine, magnesium sulfate, phenobarbital, phenobarbital sodium, phenytoin, phenytoin sodium, phenytoin sodium (extended), primidone, tiagabine hydrochloride, topiramate, valproate sodium, valproic acid. 20 The at least one antidepressant can be at least one selected from amitriptyline hydrochloride, amitriptyline pamoate, amoxapine, bupropion hydrochloride, citalopram hydrobromide, clomipramine hydrochloride, desipramine hydrochloride, doxepin hydrochloride, fluoxetine hydrochloride, imipramine hydrochloride, imipramine pamoate, mirtazapine, nefazodone hydrochloride, nortriptyline hydrochloride, paroxetine hydrochloride, phenelzine sulfate, 25 sertraline hydrochloride, tranlycypromine sulfate, trimipramine maleate, venlafaxine hydrochloride. The at least one antianxiety drug can be at least one selected from alprazolam, buspirone hydrochloride, chlordiazepoxide, chlordiazepoxide hydrochloride, clorazepate dipotassium, diazepam, doxepin hydrochloride, hydroxyzine embonate, hydroxyzine hydrochloride, hydroxyzine pamoate, lorazepam, mephrobamate, midazolam hydrochloride, 30 oxazepam. The at least one antipsychotic drug can be at least one selected from chlorpromazine hydrochloride, clozapine, fluphenazine decanoate, fluphenazine enanthate, fluphenazine hydrochloride, haloperidol, haloperidol decanoate, haloperidol lactate, loxapine hydrochloride, loxapine succinate, mesoridazine besylate, molindone hydrochloride, olanzapine, perphenazine, pimozide, prochlorperazine, quetiapine fumarate, risperidone, 35 thioridazine hydrochloride, thiothixene, thiothixene hydrochloride, trifluoperazine

hydrochloride. The at least one central nervous system stimulant can be at least one selected from amphetamine sulfate, caffeine, dextroamphetamine sulfate, doxapram hydrochloride, methamphetamine hydrochloride, methylphenidate hydrochloride, modafinil, pemoline, phentermine hydrochloride. The at least one antiparkinsonian can be at least one selected from
5 amantadine hydrochloride, benzotropine mesylate, biperiden hydrochloride, biperiden lactate, bromocriptine mesylate, carbidopa-levodopa, entacapone, levodopa, pergolide mesylate, pramipexole dihydrochloride, ropinirole hydrochloride, selegiline hydrochloride, tolcapone, trihexyphenidyl hydrochloride. The at least one miscellaneous central nervous system drug can be at least one selected from riluzole, bupropion hydrochloride, donepezil hydrochloride,
10 droperidol, fluvoxamine maleate, lithium carbonate, lithium citrate, naratriptan hydrochloride, nicotine polacrilex, nicotine transdermal system, propofol, rizatriptan benzoate, sibutramine hydrochloride monohydrate, sumatriptan succinate, tacrine hydrochloride, zolmitriptan. (See, e.g., pp. 337-530 of *Nursing 2001 Drug Handbook*.)

The at least one cholinergic (e.g., parasympathomimetic) can be at least one selected
15 from bethanechol chloride, edrophonium chloride, neostigmine bromide, neostigmine methylsulfate, physostigmine salicylate, pyridostigmine bromide. The at least one anticholinergics can be at least one selected from atropine sulfate, dicyclomine hydrochloride, glycopyrrolate, hyoscyamine, hyoscyamine sulfate, propantheline bromide, scopolamine, scopolamine butylbromide, scopolamine hydrobromide. The at least one adrenergics
20 (sympathomimetics) can be at least one selected from dobutamine hydrochloride, dopamine hydrochloride, metaraminol bitartrate, norepinephrine bitartrate, phenylephrine hydrochloride, pseudoephedrine hydrochloride, pseudoephedrine sulfate. The at least one adrenergic blocker (sympatholytic) can be at least one selected from dihydroergotamine mesylate, ergotamine tartrate, methysergide maleate, propranolol hydrochloride. The at least one skeletal muscle
25 relaxant can be at least one selected from baclofen, carisoprodol, chlorzoxazone, cyclobenzaprine hydrochloride, dantrolene sodium, methocarbamol, tizanidine hydrochloride. The at least one neuromuscular blockers can be at least one selected from atracurium besylate, cisatracurium besylate, doxacurium chloride, mivacurium chloride, pancuronium bromide, pipecuronium bromide, rapacuronium bromide, rocuronium bromide, succinylcholine chloride,
30 tubocurarine chloride, vecuronium bromide. (See, e.g., pp. 531-84 of *Nursing 2001 Drug Handbook*.)

The anti-infective drug can be at least one selected from amebicides or at least one antiprotozoals, anthelmintics, antifungals, antimalarials, antituberculotics or at least one antileprotics, aminoglycosides, penicillins, cephalosporins, tetracyclines, sulfonamides,
35 fluoroquinolones, antivirals, macrolide anti-infectives, miscellaneous anti-infectives. The CV

drug can be at least one selected from inotropics, antiarrhythmics, antianginals, antihypertensives, antilipemics, miscellaneous cardiovascular drugs. The CNS drug can be at least one selected from nonnarcotic analgesics or at least one selected from antipyretics, nonsteroidal anti-inflammatory drugs, narcotic or at least one opioid analgesics, sedative-
 5 hypnotics, anticonvulsants, antidepressants, antianxiety drugs, antipsychotics, central nervous system stimulants, antiparkinsonians, miscellaneous central nervous system drugs. The ANS drug can be at least one selected from cholinergics (parasympathomimetics), anticholinergics, adrenergics (sympathomimetics), adrenergic blockers (sympatholytics), skeletal muscle relaxants, neuromuscular blockers. The respiratory tract drug can be at least one selected from
 10 antihistamines, bronchodilators, expectorants or at least one antitussives, miscellaneous respiratory drugs. The GI tract drug can be at least one selected from antacids or at least one adsorbents or at least one antiflatulents, digestive enzymes or at least one gallstone solubilizers, antidiarrheals, laxatives, antiemetics, antiulcer drugs. The hormonal drug can be at least one selected from corticosteroids, androgens or at least one anabolic steroids, estrogens
 15 or at least one progestins, gonadotropins, antidiabetic drugs or at least one glucagon, thyroid hormones, thyroid hormone antagonists, pituitary hormones, parathyroid-like drugs. The drug for fluid and electrolyte balance can be at least one selected from diuretics, electrolytes or at least one replacement solutions, acidifiers or at least one alkalizers. The hematologic drug can be at least one selected from hematinics, anticoagulants, blood derivatives, thrombolytic
 20 enzymes. The antineoplastics can be at least one selected from alkylating drugs, antimetabolites, antibiotic antineoplastics, antineoplastics that alter hormone balance, miscellaneous antineoplastics. The immunomodulation drug can be at least one selected from immunosuppressants, vaccines or at least one toxoids, antitoxins or at least one antivenins, immune serums, biological response modifiers. The ophthalmic, otic, and nasal drugs can be
 25 at least one selected from ophthalmic anti-infectives, ophthalmic anti-inflammatories, miotics, mydriatics, ophthalmic vasoconstrictors, miscellaneous ophthalmics, otics, nasal drugs. The topical drug can be at least one selected from local anti-infectives, scabicides or at least one pediculicides, topical corticosteroids. The nutritional drug can be at least one selected from vitamins, minerals, or caloric. See, e.g., contents of *Nursing 2001 Drug Handbook, supra*.
 30 The at least one amebicide or antiprotozoal can be at least one selected from atovaquone, chloroquine hydrochloride, chloroquine phosphate, metronidazole, metronidazole hydrochloride, pentamidine isethionate. The at least one anthelmintic can be at least one selected from mebendazole, pyrantel pamoate, thiabendazole. The at least one antifungal can be at least one selected from amphotericin B, amphotericin B cholesteryl sulfate complex,
 35 amphotericin B lipid complex, amphotericin B liposomal, fluconazole, flucytosine,

griseofulvin microsize, griseofulvin ultramicrosize, itraconazole, ketoconazole, nystatin, terbinafine hydrochloride. The at least one antimalarial can be at least one selected from chloroquine hydrochloride, chloroquine phosphate, doxycycline, hydroxychloroquine sulfate, mefloquine hydrochloride, primaquine phosphate, pyrimethamine, pyrimethamine with

5 sulfadoxine. The at least one antituberculous or antileprotic can be at least one selected from clofazimine, cycloserine, dapsone, ethambutol hydrochloride, isoniazid, pyrazinamide, rifabutin, rifampin, rifapentine, streptomycin sulfate. The at least one aminoglycoside can be at least one selected from amikacin sulfate, gentamicin sulfate, neomycin sulfate, streptomycin sulfate, tobramycin sulfate. The at least one penicillin can be at least one selected from

10 amoxicillin/clavulanate potassium, amoxicillin trihydrate, ampicillin, ampicillin sodium, ampicillin trihydrate, ampicillin sodium/sulbactam sodium, cloxacillin sodium, dicloxacillin sodium, mezlocillin sodium, nafcillin sodium, oxacillin sodium, penicillin G benzathine, penicillin G potassium, penicillin G procaine, penicillin G sodium, penicillin V potassium, piperacillin sodium, piperacillin sodium/tazobactam sodium, ticarcillin disodium, ticarcillin

15 disodium/clavulanate potassium. The at least one cephalosporin can be at least one selected from at least one of cefaclor, cefadroxil, cefazolin sodium, cefdinir, cefepime hydrochloride, cefixime, cefmetazole sodium, cefonicid sodium, cefoperazone sodium, cefotaxime sodium, cefotetan disodium, cefoxitin sodium, cefpodoxime proxetil, cefprozil, ceftazidime, ceftibuten, ceftizoxime sodium, ceftriaxone sodium, cefuroxime axetil, cefuroxime sodium, cephalixin

20 hydrochloride, cephalixin monohydrate, cephradine, loracarbef. The at least one tetracycline can be at least one selected from demeclocycline hydrochloride, doxycycline calcium, doxycycline hyclate, doxycycline hydrochloride, doxycycline monohydrate, minocycline hydrochloride, tetracycline hydrochloride. The at least one sulfonamide can be at least one selected from co-trimoxazole, sulfadiazine, sulfamethoxazole, sulfisoxazole, sulfisoxazole

25 acetyl. The at least one fluoroquinolone can be at least one selected from alatrofloxacin mesylate, ciprofloxacin, enoxacin, levofloxacin, lomefloxacin hydrochloride, nalidixic acid, norfloxacin, ofloxacin, sparfloxacin, trovafloxacin mesylate. The at least one fluoroquinolone can be at least one selected from alatrofloxacin mesylate, ciprofloxacin, enoxacin, levofloxacin, lomefloxacin hydrochloride, nalidixic acid, norfloxacin, ofloxacin, sparfloxacin,

30 trovafloxacin mesylate. The at least one antiviral can be at least one selected from abacavir sulfate, acyclovir sodium, amantadine hydrochloride, amprenavir, didanosine, efavirenz, famciclovir, foscarnet sodium, foscarnet sodium, ganciclovir, indinavir sulfate, lamivudine, lamivudine/zidovudine, nelfinavir mesylate, nevirapine, oseltamivir phosphate, ribavirin, rimantadine hydrochloride, ritonavir, saquinavir,

35 saquinavir mesylate, stavudine, valacyclovir hydrochloride, zalcitabine, zanamivir, zidovudine.

- The at least one macroline anti-infective can be at least one selected from azithromycin, clarithromycin, dirithromycin, erythromycin base, erythromycin estolate, erythromycin ethylsuccinate, erythromycin lactobionate, erythromycin stearate. The at least one miscellaneous anti-infective can be at least one selected from aztreonam, bacitracin, chloramphenicol sodium succinate, clindamycin hydrochloride, clindamycin palmitate hydrochloride, clindamycin phosphate, imipenem and cilastatin sodium, meropenem, nitrofurantoin macrocrystals, nitrofurantoin microcrystals, quinupristin/dalfopristin, spectinomycin hydrochloride, trimethoprim, vancomycin hydrochloride. (See, e.g., pp. 24-214 of *Nursing 2001 Drug Handbook*.)
- The at least one inotropic can be at least one selected from amrinone lactate, digoxin, milrinone lactate. The at least one antiarrhythmic can be at least one selected from adenosine, amiodarone hydrochloride, atropine sulfate, bretylium tosylate, diltiazem hydrochloride, disopyramide, disopyramide phosphate, esmolol hydrochloride, flecainide acetate, ibutilide fumarate, lidocaine hydrochloride, mexiletine hydrochloride, moricizine hydrochloride, phenytoin, phenytoin sodium, procainamide hydrochloride, propafenone hydrochloride, propranolol hydrochloride, quinidine bisulfate, quinidine gluconate, quinidine polygalacturonate, quinidine sulfate, sotalol, tocainide hydrochloride, verapamil hydrochloride. The at least one antianginal can be at least one selected from amlodipine besylate, aml nitrite, bepridil hydrochloride, diltiazem hydrochloride, isosorbide dinitrate, isosorbide mononitrate, nadolol, nicardipine hydrochloride, nifedipine, nitroglycerin, propranolol hydrochloride, verapamil, verapamil hydrochloride. The at least one antihypertensive can be at least one selected from acebutolol hydrochloride, amlodipine besylate, atenolol, benazepril hydrochloride, betaxolol hydrochloride, bisoprolol fumarate, candesartan cilexetil, captopril, carteolol hydrochloride, carvedilol, clonidine, clonidine hydrochloride, diazoxide, diltiazem hydrochloride, doxazosin mesylate, enalaprilat, enalapril maleate, eprosartan mesylate, felodipine, fenoldopam mesylate, fosinopril sodium, guanabenz acetate, guanadrel sulfate, guanfacine hydrochloride, hydralazine hydrochloride, irbesartan, isradipine, labetalol hydrochloride, lisinopril, losartan potassium, methyldopa, methyldopate hydrochloride, metoprolol succinate, metoprolol tartrate, minoxidil, moexipril hydrochloride, nadolol, nicardipine hydrochloride, nifedipine, nisoldipine, nitroprusside sodium, penbutolol sulfate, perindopril erbumine, phentolamine mesylate, pindolol, prazosin hydrochloride, propranolol hydrochloride, quinapril hydrochloride, ramipril, telmisartan, terazosin hydrochloride, timolol maleate, trandolapril, valsartan, verapamil hydrochloride. The at least one antilipemic can be at least one selected from atorvastatin calcium, cerivastatin sodium, cholestyramine, colestipol hydrochloride, fenofibrate (micronized), fluvastatin sodium, gemfibrozil, lovastatin, niacin,

pravastatin sodium, simvastatin. The at least one miscellaneous CV drug can be at least one selected from abciximab, alprostadil, arbutamine hydrochloride, cilostazol, clopidogrel bisulfate, dipyridamole, eptifibatide, midodrine hydrochloride, pentoxifylline, ticlopidine hydrochloride, tirofiban hydrochloride. (See, e.g., pp. 215-336 of *Nursing 2001 Drug Handbook*.)

5

The at least one antihistamine can be at least one selected from brompheniramine maleate, cetirizine hydrochloride, chlorpheniramine maleate, clemastine fumarate, cyproheptadine hydrochloride, diphenhydramine hydrochloride, fexofenadine hydrochloride, loratadine, promethazine hydrochloride, promethazine theoclate, triprolidine hydrochloride.

10 The at least one bronchodilators can be at least one selected from albuterol, albuterol sulfate, aminophylline, atropine sulfate, ephedrine sulfate, epinephrine, epinephrine bitartrate, epinephrine hydrochloride, ipratropium bromide, isoproterenol, isoproterenol hydrochloride, isoproterenol sulfate, levalbuterol hydrochloride, metaproterenol sulfate, oxtriphylline, pirbuterol acetate, salmeterol xinafoate, terbutaline sulfate, theophylline. The at least one
15 expectorants or antitussives can be at least one selected from benzonatate, codeine phosphate, codeine sulfate, dextromethorphan hydrobromide, diphenhydramine hydrochloride, guaifenesin, hydromorphone hydrochloride. The at least one miscellaneous respiratory drug can be at least one selected from acetylcysteine, beclomethasone dipropionate, beractant, budesonide, calfactant, cromolyn sodium, dornase alfa, epoprostenol sodium, flunisolide,
20 fluticasone propionate, montelukast sodium, nedocromil sodium, palivizumab, triamcinolone acetonide, zafirlukast, zileuton. (See, e.g., pp. 585-642 of *Nursing 2001 Drug Handbook*.)

The at least one antacid, adsorbents, or antiflatulents can be at least one selected from aluminum carbonate, aluminum hydroxide, calcium carbonate, magaldrate, magnesium hydroxide, magnesium oxide, simethicone, sodium bicarbonate. The at least one digestive
25 enzymes or gallstone solubilizers can be at least one selected from pancreatin, pancrelipase, ursodiol. The at least one antidiarrheal can be at least one selected from attapulgite, bismuth subsalicylate, calcium polycarbophil, diphenoxylate hydrochloride or atropine sulfate, loperamide, octreotide acetate, opium tincture, opium tincture (camphorated). The at least one laxative can be at least one selected from bisocodyl, calcium polycarbophil, cascara sagrada,
30 cascara sagrada aromatic fluidextract, cascara sagrada fluidextract, castor oil, docusate calcium, docusate sodium, glycerin, lactulose, magnesium citrate, magnesium hydroxide, magnesium sulfate, methylcellulose, mineral oil, polyethylene glycol or electrolyte solution, psyllium, senna, sodium phosphates. The at least one antiemetic can be at least one selected from chlorpromazine hydrochloride, dimenhydrinate, dolasetron mesylate, dronabinol,
35 granisetron hydrochloride, meclizine hydrochloride, metoclopramide hydrochloride,

ondansetron hydrochloride, perphenazine, prochlorperazine, prochlorperazine edisylate, prochlorperazine maleate, promethazine hydrochloride, scopolamine, thiethylperazine maleate, trimethobenzamide hydrochloride. The at least one antiulcer drug can be at least one selected from cimetidine, cimetidine hydrochloride, famotidine, lansoprazole, misoprostol, nizatidine, omeprazole, rabeprazole sodium, ranitidine bismuth citrate, ranitidine hydrochloride, sucralfate. (See, e.g., pp. 643-95 of *Nursing 2001 Drug Handbook*.) The at least one corticosteroids can be at least one selected from betamethasone, betamethasone acetate or betamethasone sodium phosphate, betamethasone sodium phosphate, cortisone acetate, dexamethasone, dexamethasone acetate, dexamethasone sodium phosphate, fludrocortisone acetate, hydrocortisone, hydrocortisone acetate, hydrocortisone cypionate, hydrocortisone sodium phosphate, hydrocortisone sodium succinate, methylprednisolone, methylprednisolone acetate, methylprednisolone sodium succinate, prednisolone, prednisolone acetate, prednisolone sodium phosphate, prednisolone tebutate, prednisone, triamcinolone, triamcinolone acetonide, triamcinolone diacetate.

The at least one androgen or anabolic steroids can be at least one selected from danazol, fluoxymesterone, methyltestosterone, nandrolone decanoate, nandrolone phenpropionate, testosterone, testosterone cypionate, testosterone enanthate, testosterone propionate, testosterone transdermal system. The at least one estrogen or progestin can be at least one selected from esterified estrogens, estradiol, estradiol cypionate, estradiol/norethindrone acetate transdermal system, estradiol valerate, estrogens (conjugated), estropipate, ethinyl estradiol, ethinyl estradiol and desogestrel, ethinyl estradiol and ethynodiol diacetate, ethinyl estradiol and desogestrel, ethinyl estradiol and ethynodiol diacetate, ethinyl estradiol and levonorgestrel, ethinyl estradiol and norethindrone, ethinyl estradiol and norethindrone acetate, ethinyl estradiol and norgestimate, ethinyl estradiol and norgestrel, ethinyl estradiol and norethindrone and acetate and ferrous fumarate, levonorgestrel, medroxyprogesterone acetate, mestranol and norethindron, norethindrone, norethindrone acetate, norgestrel, progesterone. The at least one gonadotropin can be at least one selected from ganirelix acetate, gonadoreline acetate, histrelin acetate, menotropins. The at least one antidiabetic or glucaon can be at least one selected from acarbose, chlorpropamide, glimepiride, glipizide, glucagon, glyburide, insulins, metformin hydrochloride, miglitol, pioglitazone hydrochloride, repaglinide, rosiglitazone maleate, troglitazone. The at least one thyroid hormone can be at least one selected from levothyroxine sodium, liothyronine sodium, liotrix, thyroid. The at least one thyroid hormone antagonist can be at least one selected from methimazole, potassium iodide, potassium iodide (saturated solution), propylthiouracil, radioactive iodine (sodium iodide ¹³¹I), strong iodine solution. The at least one pituitary

hormone can be at least one selected from corticotropin, cosyntropin, desmopressin acetate, leuprolide acetate, repository corticotropin, somatrem, somatropin, vasopressin. The at least one parathyroid-like drug can be at least one selected from calcifediol, calcitonin (human), calcitonin (salmon), calcitriol, dihydrotachysterol, etidronate disodium. (See, e.g., pp. 696-796 of *Nursing 2001 Drug Handbook*.)

5 The at least one diuretic can be at least one selected from acetazolamide, acetazolamide sodium, amiloride hydrochloride, bumetanide, chlorthalidone, ethacrynic acid, furosemide, hydrochlorothiazide, indapamide, mannitol, metolazone, spironolactone, torsemide, triamterene, urea. The at least one electrolyte or replacement
10 solution can be at least one selected from calcium acetate, calcium carbonate, calcium chloride, calcium citrate, calcium gluconate, calcium gluceptate, calcium gluconate, calcium lactate, calcium phosphate (dibasic), calcium phosphate (tribasic), dextran (high-molecular-weight), dextran (low-molecular-weight), hetastarch, magnesium chloride, magnesium sulfate, potassium acetate, potassium bicarbonate, potassium chloride, potassium gluconate, Ringer's
15 injection, Ringer's injection (lactated), sodium chloride. The at least one acidifier or alkalizer can be at least one selected from sodium bicarbonate, sodium lactate, tromethamine. (See, e.g., pp. 797-833 of *Nursing 2001 Drug Handbook*.)

The at least one hematinic can be at least one selected from ferrous fumarate, ferrous gluconate, ferrous sulfate, ferrous sulfate (dried), iron dextran, iron sorbitol, polysaccharide-
20 iron complex, sodium ferric gluconate complex. The at least one anticoagulant can be at least one selected from ardeparin sodium, dalteparin sodium, danaparoid sodium, enoxaparin sodium, heparin calcium, heparin sodium, warfarin sodium. The at least one blood derivative can be at least one selected from albumin 5%, albumin 25%, antihemophilic factor, anti-inhibitor coagulant complex, antithrombin III (human), factor IX (human), factor IX complex,
25 plasma protein fractions. The at least one thrombolytic enzyme can be at least one selected from alteplase, anistreplase, reteplase (recombinant), streptokinase, urokinase. (See, e.g., pp. 834-66 of *Nursing 2001 Drug Handbook*.)

The at least one alkylating drug can be at least one selected from busulfan, carboplatin, carmustine, chlorambucil, cisplatin, cyclophosphamide, ifosfamide, lomustine,
30 mechlorethamine hydrochloride, melphalan, melphalan hydrochloride, streptozocin, temozolomide, thiotepa. The at least one antimetabolite can be at least one selected from capecitabine, cladribine, cytarabine, floxuridine, fludarabine phosphate, fluorouracil, hydroxyurea, mercaptopurine, methotrexate, methotrexate sodium, thioguanine. The at least one antibiotic antineoplastic can be at least one selected from bleomycin sulfate, dactinomycin,
35 daunorubicin citrate liposomal, daunorubicin hydrochloride, doxorubicin hydrochloride,

doxorubicin hydrochloride liposomal, epirubicin hydrochloride, idarubicin hydrochloride, mitomycin, pentostatin, plicamycin, valrubicin. The at least one antineoplastics that alter hormone balance can be at least one selected from anastrozole, bicalutamide, estramustine phosphate sodium, exemestane, flutamide, goserelin acetate, letrozole, leuprolide acetate, 5 megestrol acetate, nilutamide, tamoxifen citrate, testolactone, toremifene citrate. The at least one miscellaneous antineoplastic can be at least one selected from asparaginase, bacillus Calmette-Guerin (BCG) (live intravesical), dacarbazine, docetaxel, etoposide, etoposide phosphate, gemcitabine hydrochloride, irinotecan hydrochloride, mitotane, mitoxantrone hydrochloride, paclitaxel, pegaspargase, porfimer sodium, procarbazine hydrochloride, 10 rituximab, teniposide, topotecan hydrochloride, trastuzumab, tretinoin, vinblastine sulfate, vincristine sulfate, vinorelbine tartrate. (See, e.g., pp. 867-963 of *Nursing 2001 Drug Handbook*.)

The at least one immunosuppressant can be at least one selected from azathioprine, basiliximab, cyclosporine, daclizumab, lymphocyte immune globulin, muromonab-CD3, 15 mycophenolate mofetil, mycophenolate mofetil hydrochloride, sirolimus, tacrolimus. The at least one vaccine or toxoid can be at least one selected from BCG vaccine, cholera vaccine, diphtheria and tetanus toxoids (adsorbed), diphtheria and tetanus toxoids and acellular pertussis vaccine adsorbed, diphtheria and tetanus toxoids and whole-cell pertussis vaccine, *Haemophilus b* conjugate vaccines, hepatitis A vaccine (inactivated), hepatitis B vaccine 20 (recombinant), influenza virus vaccine 1999-2000 trivalent types A & B (purified surface antigen), influenza virus vaccine 1999-2000 trivalent types A & B (subvirion or purified subvirion), influenza virus vaccine 1999-2000 trivalent types A & B (whole virion), Japanese encephalitis virus vaccine (inactivated), Lyme disease vaccine (recombinant OspA), measles and mumps and rubella virus vaccine (live), measles and mumps and rubella virus vaccine (live 25 attenuated), measles virus vaccine (live attenuated), meningococcal polysaccharide vaccine, mumps virus vaccine (live), plague vaccine, pneumococcal vaccine (polyvalent), poliovirus vaccine (inactivated), poliovirus vaccine (live, oral, trivalent), rabies vaccine (adsorbed), rabies vaccine (human diploid cell), rubella and mumps virus vaccine (live), rubella virus vaccine (live, attenuated), tetanus toxoid (adsorbed), tetanus toxoid (fluid), typhoid vaccine (oral), 30 typhoid vaccine (parenteral), typhoid Vi polysaccharide vaccine, varicella virus vaccine, yellow fever vaccine. The at least one antitoxin or antivenin can be at least one selected from black widow spider antivenin, Crotalidae antivenom (polyvalent), diphtheria antitoxin (equine), *Micrurus fulvius* antivenin). The at least one immune serum can be at least one selected from cytomegalovirus immune globulin (intravenous), hepatitis B immune globulin 35 (human), immune globulin intramuscular, immune globulin intravenous, rabies immune

globulin (human), respiratory syncytial virus immune globulin intravenous (human), Rh₀(D) immune globulin (human), Rh₀(D) immune globulin intravenous (human), tetanus immune globulin (human), varicella-zoster immune globulin. The at least one biological response modifiers can be at least one selected from aldesleukin, epoetin alfa, filgrastim, glatiramer acetate for injection, interferon alfacon-1, interferon alfa-2a (recombinant), interferon alfa-2b (recombinant), interferon beta-1a, interferon beta-1b (recombinant), interferon gamma-1b, levamisole hydrochloride, oprelvekin, sargramostim. (See, e.g., pp. 964-1040 of *Nursing 2001 Drug Handbook*.)

The at least one ophthalmic anti-infectives can be selected from bacitracin, chloramphenicol, ciprofloxacin hydrochloride, erythromycin, gentamicin sulfate, ofloxacin 0.3%, polymyxin B sulfate, sulfacetamide sodium 10%, sulfacetamide sodium 15%, sulfacetamide sodium 30%, tobramycin, vidarabine. The at least one ophthalmic anti-inflammatories can be at least one selected from dexamethasone, dexamethasone sodium phosphate, diclofenac sodium 0.1%, fluorometholone, flurbiprofen sodium, ketorolac tromethamine, prednisolone acetate (suspension) prednisolone sodium phosphate (solution). The at least one miotic can be at least one selected from acetylcholine chloride, carbachol (intraocular), carbachol (topical), echothiophate iodide, pilocarpine, pilocarpine hydrochloride, pilocarpine nitrate. The at least one mydriatic can be at least one selected from atropine sulfate, cyclopentolate hydrochloride, epinephrine hydrochloride, epinephryl borate, homatropine hydrobromide, phenylephrine hydrochloride, scopolamine hydrobromide, tropicamide. The at least one ophthalmic vasoconstrictors can be at least one selected from naphazoline hydrochloride, oxymetazoline hydrochloride, tetrahydrozoline hydrochloride. The at least one miscellaneous ophthalmics can be at least one selected from apraclonidine hydrochloride, betaxolol hydrochloride, brimonidine tartrate, carteolol hydrochloride, dipivefrin hydrochloride, dorzolamide hydrochloride, emedastine difumarate, fluorescein sodium, ketotifen fumarate, latanoprost, levobunolol hydrochloride, metipranolol hydrochloride, sodium chloride (hypertonic), timolol maleate. The at least one otic can be at least one selected from boric acid, carbamide peroxide, chloramphenicol, triethanolamine polypeptide oleate-condensate. The at least one nasal drug can be at least one selected from beclomethasone dipropionate, budesonide, ephedrine sulfate, epinephrine hydrochloride, flunisolide, fluticasone propionate, naphazoline hydrochloride, oxymetazoline hydrochloride, phenylephrine hydrochloride, tetrahydrozoline hydrochloride, triamcinolone acetonide, xylometazoline hydrochloride. (See, e.g., pp. 1041-97 of *Nursing 2001 Drug Handbook*.)

The at least one local anti-infectives can be at least one selected from acyclovir, amphotericin B, azelaic acid cream, bacitracin, butoconazole nitrate, clindamycin phosphate,

clotrimazole, econazole nitrate, erythromycin, gentamicin sulfate, ketoconazole, mafenide acetate, metronidazole (topical), miconazole nitrate, mupirocin, naftifine hydrochloride, neomycin sulfate, nitrofurazone, nystatin, silver sulfadiazine, terbinafine hydrochloride, terconazole, tetracycline hydrochloride, tioconazole, tolnaftate. The at least one scabicide or
5 pediculicide can be at least one selected from crotamiton, lindane, permethrin, pyrethrins. The at least one topical corticosteroid can be at least one selected from betamethasone dipropionate, betamethasone valerate, clobetasol propionate, desonide, desoximetasone, dexamethasone, dexamethasone sodium phosphate, diflorasone diacetate, fluocinolone acetonide, fluocinonide, flurandrenolide, fluticasone propionate, halcionide, hydrocortisone,
10 hydrocortisone acetate, hydrocortisone butyrate, hydrocortisone valerate, mometasone furoate, triamcinolone acetonide. (See, e.g., pp. 1098-1136 of *Nursing 2001 Drug Handbook*.)

The at least one vitamin or mineral can be at least one selected from vitamin A, vitamin B complex, cyanocobalamin, folic acid, hydroxocobalamin, leucovorin calcium, niacin, niacinamide, pyridoxine hydrochloride, riboflavin, thiamine hydrochloride, vitamin C,
15 vitamin D, cholecalciferol, ergocalciferol, vitamin D analogue, doxercalciferol, paricalcitol, vitamin E, vitamin K analogue, phytonadione, sodium fluoride, sodium fluoride (topical), trace elements, chromium, copper, iodine, manganese, selenium, zinc. The at least one calorics can be at least one selected from amino acid infusions (crystalline), amino acid infusions in dextrose, amino acid infusions with electrolytes, amino acid infusions with electrolytes in
20 dextrose, amino acid infusions for hepatic failure, amino acid infusions for high metabolic stress, amino acid infusions for renal failure, dextrose, fat emulsions, medium-chain triglycerides. (See, e.g., pp. 1137-63 of *Nursing 2001 Drug Handbook*.)

Anti-amyloid antibody compositions of the present invention can further comprise at least one of any suitable and effective amount of a composition or pharmaceutical composition
25 comprising at least one anti-amyloid antibody to a cell, tissue, organ, animal or patient in need of such modulation, treatment or therapy, optionally further comprising at least one selected from at least one TNF antagonist (e.g., but not limited to a TNF chemical or protein antagonist, TNF monoclonal or polyclonal antibody or fragment, a soluble TNF receptor (e.g., p55, p70 or p85) or fragment, fusion polypeptides thereof, or a small molecule TNF antagonist, e.g., TNF
30 binding protein I or II (TBP-I or TBP-II), nerelimonmab, infliximab, entercept, CDP-571, CDP-870, afelimomab, lenercept, and the like), an antirheumatic (e.g., methotrexate, auranofin, aurothioglucose, azathioprine, etanercept, gold sodium thiomalate, hydroxychloroquine sulfate, leflunomide, sulfasalazine), a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an
35 antimicrobial (e.g., aminoglycoside, an antifungal, an antiparasitic, an antiviral, a carbapenem,

cephalosporin, a fluorquinolone, a macrolide, a penicillin, a sulfonamide, a tetracycline, another antimicrobial), an antipsoriatic, a corticosteroid, an anabolic steroid, a diabetes related agent, a mineral, a nutritional, a thyroid agent, a vitamin, a calcium related hormone, an antidiarrheal, an antitussive, an antiemetic, an antiulcer, a laxative, an anticoagulant, an erythropoietin (e.g., epoetin alpha), a filgrastim (e.g., G-CSF, Neupogen), a sargramostim (GM-CSF, Leukine), an immunization, an immunoglobulin, an immunosuppressive (e.g., basiliximab, cyclosporine, daclizumab), a growth hormone, a hormone replacement drug, an estrogen receptor modulator, a mydriatic, a cycloplegic, an alkylating agent, an antimetabolite, a mitotic inhibitor, a radiopharmaceutical, an antidepressant, antimanic agent, an antipsychotic, an anxiolytic, a hypnotic, a sympathomimetic, a stimulant, donepezil, tacrine, an asthma medication, a beta agonist, an inhaled steroid, a leukotriene inhibitor, a methylxanthine, a cromolyn, an epinephrine or analog, dornase alpha (Pulmozyme), a cytokine or a cytokine antagonist. Non-limiting examples of such cytokines include, but are not limited to, any of IL-1 to IL-23. Suitable dosages are well known in the art. See, e.g., Wells et al., eds., *Pharmacotherapy Handbook*, 2nd Edition, Appleton and Lange, Stamford, CT (2000); *PDR Pharmacopoeia*, Tarascon Pocket Pharmacopoeia 2000, Deluxe Edition, Tarascon Publishing, Loma Linda, CA (2000), each of which references are entirely incorporated herein by reference.

Such anti-cancer or anti-infectives can also include toxin molecules that are associated, bound, co-formulated or co-administered with at least one antibody of the present invention. The toxin can optionally act to selectively kill the pathologic cell or tissue. The pathologic cell can be a cancer or other cell. Such toxins can be, but are not limited to, purified or recombinant toxin or toxin fragment comprising at least one functional cytotoxic domain of toxin, e.g., selected from at least one of ricin, diphtheria toxin, a venom toxin, or a bacterial toxin. The term toxin also includes both endotoxins and exotoxins produced by any naturally occurring, mutant or recombinant bacteria or viruses which may cause any pathological condition in humans and other mammals, including toxin shock, which can result in death. Such toxins may include, but are not limited to, enterotoxigenic *E. coli* heat-labile enterotoxin (LT), heat-stable enterotoxin (ST), *Shigella* cytotoxin, *Aeromonas* enterotoxins, toxic shock syndrome toxin-1 (TSST-1), Staphylococcal enterotoxin A (SEA), B (SEB), or C (SEC), Streptococcal enterotoxins and the like. Such bacteria include, but are not limited to, strains of a species of enterotoxigenic *E. coli* (ETEC), enterohemorrhagic *E. coli* (e.g., strains of serotype O157:H7), Staphylococcus species (e.g., *Staphylococcus aureus*, *Staphylococcus pyogenes*), *Shigella* species (e.g., *Shigella dysenteriae*, *Shigella flexneri*, *Shigella boydii*, and *Shigella sonnei*), *Salmonella* species (e.g., *Salmonella typhi*, *Salmonella cholerae-suis*, *Salmonella*

enteritidis), *Clostridium* species (e.g., *Clostridium perfringens*, *Clostridium difficile*, *Clostridium botulinum*), *Camphlobacter* species (e.g., *Camphlobacter jejuni*, *Camphlobacter fetus*), *Heliobacter* species, (e.g., *Heliobacter pylori*), *Aeromonas* species (e.g., *Aeromonas sobria*, *Aeromonas hydrophila*, *Aeromonas caviae*), *Pleisomonas shigelloides*, *Yersina enterocolitica*, *Vibrios* species (e.g., *Vibrios cholerae*, *Vibrios parahemolyticus*), *Klebsiella* species, *Pseudomonas aeruginosa*, and *Streptococci*. See, e.g., Stein, ed., INTERNAL MEDICINE, 3rd ed., pp 1-13, Little, Brown and Co., Boston, (1990); Evans et al., eds., Bacterial Infections of Humans: Epidemiology and Control, 2d. Ed., pp 239-254, Plenum Medical Book Co., New York (1991); Mandell et al, Principles and Practice of Infectious Diseases, 3d. Ed., Churchill Livingstone, New York (1990); Berkow et al, eds., *The Merck Manual*, 16th edition, Merck and Co., Rahway, N.J., 1992; Wood et al, FEMS Microbiology Immunology, 76:121-134 (1991); Marrack et al, Science, 248:705-711 (1990), the contents of which references are incorporated entirely herein by reference.

Anti-amyloid antibody compounds, compositions or combinations of the present invention can further comprise at least one of any suitable auxiliary, such as, but not limited to, diluent, binder, stabilizer, buffers, salts, lipophilic solvents, preservative, adjuvant or the like. Pharmaceutically acceptable auxiliaries are preferred. Non-limiting examples of, and methods of preparing such sterile solutions are well known in the art, such as, but limited to, Gennaro, Ed., *Remington's Pharmaceutical Sciences*, 18th Edition, Mack Publishing Co. (Easton, PA) 1990. Pharmaceutically acceptable carriers can be routinely selected that are suitable for the mode of administration, solubility and/or stability of the anti-amyloid antibody, fragment or variant composition as well known in the art or as described herein.

Pharmaceutical excipients and additives useful in the present composition include but are not limited to proteins, peptides, amino acids, lipids, and carbohydrates (e.g., sugars, including monosaccharides, di-, tri-, tetra-, and oligosaccharides; derivatized sugars such as alditols, aldonic acids, esterified sugars and the like; and polysaccharides or sugar polymers), which can be present singly or in combination, comprising alone or in combination 1-99.99% by weight or volume. Exemplary protein excipients include serum albumin such as human serum albumin (HSA), recombinant human albumin (rHA), gelatin, casein, and the like. Representative amino acid/antibody components, which can also function in a buffering capacity, include alanine, glycine, arginine, betaine, histidine, glutamic acid, aspartic acid, cysteine, lysine, leucine, isoleucine, valine, methionine, phenylalanine, aspartame, and the like. One preferred amino acid is glycine.

Carbohydrate excipients suitable for use in the invention include, for example,

monosaccharides such as fructose, maltose, galactose, glucose, D-mannose, sorbose, and the like; disaccharides, such as lactose, sucrose, trehalose, cellobiose, and the like; polysaccharides, such as raffinose, melezitose, maltodextrins, dextrans, starches, and the like; and alditols, such as mannitol, xylitol, maltitol, lactitol, xylitol sorbitol (glucitol), myoinositol and the like. Preferred carbohydrate excipients for use in the present invention are mannitol, trehalose, and raffinose.

Anti-amyloid antibody compositions can also include a buffer or a pH adjusting agent; typically, the buffer is a salt prepared from an organic acid or base. Representative buffers include organic acid salts such as salts of citric acid, ascorbic acid, gluconic acid, carbonic acid, tartaric acid, succinic acid, acetic acid, or phthalic acid; Tris, tromethamine hydrochloride, or phosphate buffers. Preferred buffers for use in the present compositions are organic acid salts such as citrate.

Additionally, anti-amyloid antibody compositions of the invention can include polymeric excipients/additives such as polyvinylpyrrolidones, ficolls (a polymeric sugar), dextrans (e.g., cyclodextrins, such as 2-hydroxypropyl- β -cyclodextrin), polyethylene glycols, flavoring agents, antimicrobial agents, sweeteners, antioxidants, antistatic agents, surfactants (e.g., polysorbates such as "TWEEN 20" and "TWEEN 80"), lipids (e.g., phospholipids, fatty acids), steroids (e.g., cholesterol), and chelating agents (e.g., EDTA).

These and additional known pharmaceutical excipients and/or additives suitable for use in the anti-amyloid antibody, portion or variant compositions according to the invention are known in the art, e.g., as listed in "Remington: The Science & Practice of Pharmacy", 19th ed., Williams & Williams, (1995), and in the "Physician's Desk Reference", 52nd ed., Medical Economics, Montvale, NJ (1998), the disclosures of which are entirely incorporated herein by reference. Preferred carrier or excipient materials are carbohydrates (e.g., saccharides and alditols) and buffers (e.g., citrate) or polymeric agents.

Formulations

As noted above, the invention provides for stable formulations, which is preferably a phosphate buffer with saline or a chosen salt, as well as preserved solutions and formulations containing a preservative as well as multi-use preserved formulations suitable for pharmaceutical or veterinary use, comprising at least one anti-amyloid antibody in a pharmaceutically acceptable formulation. Preserved formulations contain at least one known preservative or optionally selected from the group consisting of at least one phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, phenylmercuric nitrite, phenoxyethanol, formaldehyde, chlorobutanol, magnesium chloride (e.g., hexahydrate), alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium

dehydroacetate and thimerosal, or mixtures thereof in an aqueous diluent. Any suitable concentration or mixture can be used as known in the art, such as 0.001-5%, or any range or value therein, such as, but not limited to 0.001, 0.003, 0.005, 0.009, 0.01, 0.02, 0.03, 0.05, 0.09, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.3, 4.5, 4.6, 4.7, 4.8, 4.9, or any range or value therein. Non-limiting examples include, no preservative, 0.1-2% m-cresol (e.g., 0.2, 0.3, 0.4, 0.5, 0.9, 1.0%), 0.1-3% benzyl alcohol (e.g., 0.5, 0.9, 1.1, 1.5, 1.9, 2.0, 2.5%), 0.001-0.5% thimerosal (e.g., 0.005, 0.01), 0.001-2.0% phenol (e.g., 0.05, 0.25, 0.28, 0.5, 0.9, 1.0%), 0.0005-1.0% alkylparaben(s) (e.g., 0.00075, 0.0009, 0.001, 0.002, 0.005, 0.0075, 0.009, 0.01, 0.02, 0.05, 0.075, 0.09, 0.1, 0.2, 0.3, 0.5, 0.75, 0.9, 1.0%), and the like.

As noted above, the invention provides an article of manufacture, comprising packaging material and at least one vial comprising a solution of at least one anti-amyloid antibody with the prescribed buffers and/or preservatives, optionally in an aqueous diluent, wherein said packaging material comprises a label that indicates that such solution can be held over a period of 1, 2, 3, 4, 5, 6, 9, 12, 18, 20, 24, 30, 36, 40, 48, 54, 60, 66, 72 hours or greater. The invention further comprises an article of manufacture, comprising packaging material, a first vial comprising lyophilized at least one anti-amyloid antibody, and a second vial comprising an aqueous diluent of prescribed buffer or preservative, wherein said packaging material comprises a label that instructs a patient to reconstitute the at least one anti-amyloid antibody in the aqueous diluent to form a solution that can be held over a period of twenty-four hours or greater.

The at least one anti-amyloid antibody used in accordance with the present invention can be produced by recombinant means, including from mammalian cell or transgenic preparations, or can be purified from other biological sources, as described herein or as known in the art.

The range of at least one anti-amyloid antibody in the product of the present invention includes amounts yielding upon reconstitution, if in a wet/dry system, concentrations from about 1.0 µg/ml to about 1000 mg/ml, although lower and higher concentrations are operable and are dependent on the intended delivery vehicle, e.g., solution formulations will differ from transdermal patch, pulmonary, transmucosal, or osmotic or micro pump methods.

Preferably, the aqueous diluent optionally further comprises a pharmaceutically acceptable preservative. Preferred preservatives include those selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride,

sodium dehydroacetate and thimerosal, or mixtures thereof. The concentration of preservative used in the formulation is a concentration sufficient to yield an anti-microbial effect. Such concentrations are dependent on the preservative selected and are readily determined by the skilled artisan.

5 Other excipients, e.g. isotonicity agents, buffers, antioxidants, preservative enhancers, can be optionally and preferably added to the diluent. An isotonicity agent, such as glycerin, is commonly used at known concentrations. A physiologically tolerated buffer is preferably added to provide improved pH control. The formulations can cover a wide range of pHs, such as from about pH 4 to about pH 10, and preferred ranges from about pH 5 to about pH 9, and a
10 most preferred range of about 6.0 to about 8.0. Preferably the formulations of the present invention have pH between about 6.8 and about 7.8. Preferred buffers include phosphate buffers, most preferably sodium phosphate, particularly phosphate buffered saline (PBS).

 Other additives, such as a pharmaceutically acceptable solubilizers like Tween
20 (polyoxyethylene (20) sorbitan monolaurate), Tween 40 (polyoxyethylene (20) sorbitan
15 monopalmitate), Tween 80 (polyoxyethylene (20) sorbitan monooleate), Pluronic F68 (polyoxyethylene polyoxypropylene block copolymers), and PEG (polyethylene glycol) or non-ionic surfactants such as polysorbate 20 or 80 or poloxamer 184 or 188, Pluronic® polyols, other block co-polymers, and chelators such as EDTA and EGTA can optionally be added to the formulations or compositions to reduce aggregation. These additives are particularly useful
20 if a pump or plastic container is used to administer the formulation. The presence of pharmaceutically acceptable surfactant mitigates the propensity for the protein to aggregate.

 The formulations of the present invention can be prepared by a process which comprises mixing at least one anti-amyloid antibody and a preservative selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben,
25 (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal or mixtures thereof in an aqueous diluent. Mixing the at least one anti-amyloid antibody and preservative in an aqueous diluent is carried out using conventional dissolution and mixing procedures. To prepare a suitable formulation, for example, a measured amount of at least one anti-amyloid antibody in buffered solution is
30 combined with the desired preservative in a buffered solution in quantities sufficient to provide the protein and preservative at the desired concentrations. Variations of this process would be recognized by one of ordinary skill in the art. For example, the order the components are added, whether additional additives are used, the temperature and pH at which the formulation is prepared, are all factors that can be optimized for the concentration and means of
35 administration used.

The claimed formulations can be provided to patients as clear solutions or as dual vials comprising a vial of lyophilized at least one anti-amyloid antibody that is reconstituted with a second vial containing water, a preservative and/or excipients, preferably a phosphate buffer and/or saline and a chosen salt, in an aqueous diluent. Either a single solution vial or dual vial requiring reconstitution can be reused multiple times and can suffice for a single or multiple cycles of patient treatment and thus can provide a more convenient treatment regimen than currently available.

The present claimed articles of manufacture are useful for administration over a period of immediately to twenty-four hours or greater. Accordingly, the presently claimed articles of manufacture offer significant advantages to the patient. Formulations of the invention can optionally be safely stored at temperatures of from about 2 to about 40°C and retain the biological activity of the protein for extended periods of time, thus, allowing a package label indicating that the solution can be held and/or used over a period of 6, 12, 18, 24, 36, 48, 72, or 96 hours or greater. If preserved diluent is used, such label can include use up to 1-12 months, one-half, one and a half, and/or two years.

The solutions of at least one anti-amyloid antibody in the invention can be prepared by a process that comprises mixing at least one antibody in an aqueous diluent. Mixing is carried out using conventional dissolution and mixing procedures. To prepare a suitable diluent, for example, a measured amount of at least one antibody in water or buffer is combined in quantities sufficient to provide the protein and optionally a preservative or buffer at the desired concentrations. Variations of this process would be recognized by one of ordinary skill in the art. For example, the order the components are added, whether additional additives are used, the temperature and pH at which the formulation is prepared, are all factors that can be optimized for the concentration and means of administration used.

The claimed products can be provided to patients as clear solutions or as dual vials comprising a vial of lyophilized at least one anti-amyloid antibody that is reconstituted with a second vial containing the aqueous diluent. Either a single solution vial or dual vial requiring reconstitution can be reused multiple times and can suffice for a single or multiple cycles of patient treatment and thus provides a more convenient treatment regimen than currently available.

The claimed products can be provided indirectly to patients by providing to pharmacies, clinics, or other such institutions and facilities, clear solutions or dual vials comprising a vial of lyophilized at least one anti-amyloid antibody that is reconstituted with a second vial containing the aqueous diluent. The clear solution in this case can be up to one liter or even larger in size, providing a large reservoir from which smaller portions of the at

least one antibody solution can be retrieved one or multiple times for transfer into smaller vials and provided by the pharmacy or clinic to their customers and/or patients.

Recognized devices comprising these single vial systems include those pen-injector devices for delivery of a solution such as BD Pens, BD Autojector[®], Humaject[®],
5 NovoPen[®], B-D[®]Pen, AutoPen[®], and OptiPen[®], GenotropinPen[®], Genotronorm Pen[®], Humatro
Pen[®], Reco-Pen[®], Roferon Pen[®], Biojector[®], Iject[®], J-tip Needle-Free Injector[®], Intraject[®],
Medi-Ject[®], e.g., as made or developed by Becton Dickenson (Franklin Lakes, NJ,
www.bectondickenson.com), Disetronic (Burgdorf, Switzerland, www.disetronic.com; Bioject,
Portland, Oregon (www.bioject.com); National Medical Products , Weston Medical
10 (Peterborough, UK, www.weston-medical.com), Medi-Ject Corp (Minneapolis, MN,
www.mediject.com). Recognized devices comprising a dual vial system include those pen-
injector systems for reconstituting a lyophilized drug in a cartridge for delivery of the
reconstituted solution such as the HumatroPen[®].

The products presently claimed include packaging material. The packaging
15 material provides, in addition to the information required by the regulatory agencies, the
conditions under which the product can be used. The packaging material of the present
invention provides instructions to the patient to reconstitute the at least one anti-amyloid
antibody in the aqueous diluent to form a solution and to use the solution over a period of 2-24
hours or greater for the two vial, wet/dry, product. For the single vial, solution product, the
20 label indicates that such solution can be used over a period of 2-24 hours or greater. The
presently claimed products are useful for human pharmaceutical product use.

The formulations of the present invention can be prepared by a process that
comprises mixing at least one anti-amyloid antibody and a selected buffer, preferably a
phosphate buffer containing saline or a chosen salt. Mixing the at least one anti-amyloid
25 antibody and buffer in an aqueous diluent is carried out using conventional dissolution and
mixing procedures. To prepare a suitable formulation, for example, a measured amount of at
least one antibody in water or buffer is combined with the desired buffering agent in water in
quantities sufficient to provide the protein and buffer at the desired concentrations. Variations
of this process would be recognized by one of ordinary skill in the art. For example, the order
30 the components are added, whether additional additives are used, the temperature and pH at
which the formulation is prepared, are all factors that can be optimized for the concentration
and means of administration used.

The claimed stable or preserved formulations can be provided to patients as
clear solutions or as dual vials comprising a vial of lyophilized at least one anti-amyloid
35 antibody that is reconstituted with a second vial containing a preservative or buffer and

excipients in an aqueous diluent. Either a single solution vial or dual vial requiring reconstitution can be reused multiple times and can suffice for a single or multiple cycles of patient treatment and thus provides a more convenient treatment regimen than currently available.

5 Other formulations or methods of stabilizing the anti-amyloid antibody may result in other than a clear solution of lyophilized powder comprising said antibody. Among non-clear solutions are formulations comprising particulate suspensions, said particulates being a composition containing the anti-amyloid antibody in a structure of variable dimension and known variously as a microsphere, microparticle, nanoparticle, nanosphere, or liposome.

10 Such relatively homogenous essentially spherical particulate formulations containing an active agent can be formed by contacting an aqueous phase containing the active and a polymer and a nonaqueous phase followed by evaporation of the nonaqueous phase to cause the coalescence of particles from the aqueous phase as taught in U.S. 4,589,330. Porous microparticles can be prepared using a first phase containing active and a polymer dispersed in a continuous solvent

15 and removing said solvent from the suspension by freeze-drying or dilution-extraction-precipitation as taught in U.S. 4,818,542. Preferred polymers for such preparations are natural or synthetic copolymers or polymer selected from the group consisting of gellan, agar, starch, arabinogalactan, albumin, collagen, polyglycolic acid, polylactic acid, glycolide-L(-) lactide poly(epsilon-caprolactone, poly(epsilon-caprolactone-CO-lactic acid), poly(epsilon-caprolactone-CO-glycolic acid), poly(beta-hydroxy butyric acid), polyethylene oxide,

20 polyethylene, poly(alkyl-2-cyanoacrylate), poly(hydroxyethyl methacrylate), polyamides, poly(amino acids), poly(2-hydroxyethyl DL-aspartamide), poly(ester urea), poly(L-phenylalanine/ethylene glycol/1,6-diisocyanatohexane) and poly(methyl methacrylate). Particularly preferred polymers are polyesters such as polyglycolic acid, polylactic acid,

25 glycolide-L(-) lactide poly(epsilon-caprolactone, poly(epsilon-caprolactone-CO-lactic acid), and poly(epsilon-caprolactone-CO-glycolic acid. Solvents useful for dissolving the polymer and/or the active include: water, hexafluoroisopropanol, methylenechloride, tetrahydrofuran, hexane, benzene, or hexafluoroacetone sesquihydrate. The process of dispersing the active containing phase with a second phase may include pressure forcing said first phase through an

30 orifice in a nozzle to affect droplet formation.

 Dry powder formulations may result from processes other than lyophilization such as by spray drying or solvent extraction by evaporation or by precipitation of a crystalline composition followed by one or more steps to remove aqueous or nonaqueous solvent. Preparation of a spray-dried antibody preparation is taught in U.S. 6,019,968. The antibody-

35 based dry powder compositions may be produced by spray drying solutions or slurries of the

antibody and, optionally, excipients, in a solvent under conditions to provide a respirable dry powder. Solvents may include polar compounds such as water and ethanol, which may be readily dried. Antibody stability may be enhanced by performing the spray drying procedures in the absence of oxygen, such as under a nitrogen blanket or by using nitrogen as the drying gas. Another relatively dry formulation is a dispersion of a plurality of perforated microstructures dispersed in a suspension medium that typically comprises a hydrofluoroalkane propellant as taught in WO 9916419. The stabilized dispersions may be administered to the lung of a patient using a metered dose inhaler. Equipment useful in the commercial manufacture of spray dried medicaments are manufactured by Buchi Ltd. or Niro Corp.

At least one anti-amyloid antibody in either the stable or preserved formulations or solutions described herein, can be administered to a patient in accordance with the present invention via a variety of delivery methods including SC or IM injection; transdermal, pulmonary, transmucosal, implant, osmotic pump, cartridge, micro pump, or other means appreciated by the skilled artisan, as well-known in the art.

Therapeutic Applications

The present invention also provides a method for modulating or treating at least one amyloid related disease, in a cell, tissue, organ, animal, or patient, as known in the art or as described herein, using at least one amyloid antibody of the present invention.

The present invention also provides a method for modulating or treating at least one amyloid related disease, in a cell, tissue, organ, animal, or patient including, but not limited to, at least one of obesity, an immune related disease, a cardiovascular disease, an infectious disease, a malignant disease or a neurologic disease. Such amyloid related diseases can include, but are not limited to, any amyloidosis, systemic amyloidosis, Alzheimer's disease (AD), sporadic Alzheimer's disease, familial Alzheimer's disease, Lewy body variant Alzheimer's disease, prion diseases, primary systemic amyloidosis, secondary systemic amyloidosis, dense systemic amyloidosis, monoclonal protein systemic amyloidosis, reactive systemic amyloidosis, hereditary apoA1 amyloidosis, hereditary lysozyme amyloidosis, insulin related amyloid, familial amyloidosis Finnish type, familial subepithelial corneal amyloid, familial amyloid polyneuropathy, familial non-neuropathic amyloidosis, familial British dementia, hereditary cerebral amyloid angiopathy, hemodialysis related amyloidosis, familial amyloid polyneuropathy, familial amyloidotic polyneuropathy, maturity onset diabetes, type II diabetes, hereditary renal amyloidosis, pituitary gland amyloidosis, injection-localization amyloidosis, medullary carcinoma, medullary carcinoma of the thyroid, atrial amyloidosis, isolated atrial amyloidosis, hereditary cerebral amyloid angiopathy, hereditary fibrinogen

orchitis/vasectomy reversal procedures, allergic/atopic diseases, asthma, allergic rhinitis, eczema, allergic contact dermatitis, allergic conjunctivitis, hypersensitivity pneumonitis, transplants, organ transplant rejection, graft-versus-host disease, systemic inflammatory response syndrome, sepsis syndrome, gram positive sepsis, gram negative sepsis, culture
 5 negative sepsis, fungal sepsis, neutropenic fever, urosepsis, meningococemia, trauma/hemorrhage, burns, ionizing radiation exposure, acute pancreatitis, adult respiratory distress syndrome, rheumatoid arthritis, alcohol-induced hepatitis, chronic inflammatory pathologies, sarcoidosis, Crohn's pathology, sickle cell anemia, diabetes, nephrosis, atopic diseases, hypersensitivity reactions, allergic rhinitis, hay fever, perennial rhinitis, conjunctivitis,
 10 endometriosis, asthma, urticaria, systemic anaphalaxis, dermatitis, pernicious anemia, hemolytic disease, thrombocytopenia, graft rejection of any organ or tissue, kidney transplant rejection, heart transplant rejection, liver transplant rejection, pancreas transplant rejection, lung transplant rejection, bone marrow transplant (BMT) rejection, skin allograft rejection, cartilage transplant rejection, bone graft rejection, small bowel transplant rejection,
 15 fetal thymus implant rejection, parathyroid transplant rejection, xenograft rejection of any organ or tissue, allograft rejection, anti-receptor hypersensitivity reactions, Graves disease, Raynaud's disease, type B insulin-resistant diabetes, asthma, myasthenia gravis, antibody-mediated cytotoxicity, type III hypersensitivity reactions, systemic lupus erythematosus, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal gammopathy, and skin changes syndrome), polyneuropathy, organomegaly, endocrinopathy, monoclonal
 20 gammopathy, skin changes syndrome, antiphospholipid syndrome, pemphigus, scleroderma, mixed connective tissue disease, idiopathic Addison's disease, diabetes mellitus, chronic active hepatitis, primary biliary cirrhosis, vitiligo, vasculitis, post-MI cardiomyopathy syndrome, type IV hypersensitivity, contact dermatitis, hypersensitivity pneumonitis, allograft rejection,
 25 granulomas due to intracellular organisms, drug sensitivity, metabolic/idiopathic, Wilson's disease, hemochromatosis, alpha-1-antitrypsin deficiency, diabetic retinopathy, hashimoto's thyroiditis, osteoporosis, hypothalamic-pituitary-adrenal axis evaluation, primary biliary cirrhosis, thyroiditis, encephalomyelitis, cachexia, cystic fibrosis, neonatal chronic lung disease, chronic obstructive pulmonary disease (COPD), familial hemophagocytic
 30 lymphohistiocytosis, dermatologic conditions, psoriasis, alopecia, nephrotic syndrome, nephritis, glomerular nephritis, acute renal failure, hemodialysis, uremia, toxicity, preeclampsia, okt3 therapy, anti-cd3 therapy, cytokine therapy, chemotherapy, radiation therapy (e.g., including but not limited to anemias, cachexia, and the like), chronic salicylate intoxication, and the like. See, e.g., the Merck Manual, 12th-17th Editions, Merck
 35 & Company, Rahway, NJ (1972, 1977, 1982, 1987, 1992, 1999), Pharmacotherapy Handbook,

Wells et al., eds., Second Edition, Appleton and Lange, Stamford, Conn. (1998, 2000), each entirely incorporated by reference.

The present invention also provides a method for modulating or treating at least one cardiovascular or amyloid related disease in a cell, tissue, organ, animal, or patient, including, but not limited to, at least one of cardiac stun syndrome, myocardial infarction, congestive heart failure, stroke, ischemic stroke, hemorrhage, arteriosclerosis, atherosclerosis, restenosis, diabetic atherosclerotic disease, hypertension, arterial hypertension, renovascular hypertension, syncope, shock, syphilis of the cardiovascular system, heart failure, cor pulmonale, primary pulmonary hypertension, cardiac arrhythmias, atrial ectopic beats, atrial flutter, atrial fibrillation (sustained or paroxysmal), post perfusion syndrome, cardiopulmonary bypass inflammation response, chaotic or multifocal atrial tachycardia, regular narrow QRS tachycardia, specific arrhythmias, ventricular fibrillation, His bundle arrhythmias, atrioventricular block, bundle branch block, myocardial ischemic disorders, coronary artery disease, angina pectoris, myocardial infarction, cardiomyopathy, dilated congestive cardiomyopathy, restrictive cardiomyopathy, valvular heart diseases, endocarditis, pericardial disease, cardiac tumors, aortic and peripheral aneurysms, aortic dissection, inflammation of the aorta, occlusion of the abdominal aorta and its branches, peripheral vascular disorders, occlusive arterial disorders, peripheral atherosclerotic disease, thromboangitis obliterans, functional peripheral arterial disorders, Raynaud's phenomenon and disease, acrocyanosis, erythromelalgia, venous diseases, venous thrombosis, varicose veins, arteriovenous fistula, lymphedema, lipedema, unstable angina, reperfusion injury, post pump syndrome, ischemia-reperfusion injury, and the like. Such a method can optionally comprise administering an effective amount of a composition or pharmaceutical composition comprising at least one anti-amyloid antibody to a cell, tissue, organ, animal or patient in need of such modulation, treatment or therapy.

The present invention also provides a method for modulating or treating at least one infectious or amyloid related disease in a cell, tissue, organ, animal or patient, including, but not limited to, at least one of: acute or chronic bacterial infection, acute and chronic parasitic or infectious processes, including bacterial, viral and fungal infections, HIV infection/HIV neuropathy, meningitis, hepatitis (e.g., A,B or C, or the like), septic arthritis, peritonitis, pneumonia, epiglottitis, e. coli 0157:h7, hemolytic uremic syndrome/thrombolytic thrombocytopenic purpura, malaria, dengue hemorrhagic fever, leishmaniasis, leprosy, toxic shock syndrome, streptococcal myositis, gas gangrene, mycobacterium tuberculosis, mycobacterium avium intracellulare, pneumocystis carinii pneumonia, pelvic inflammatory

disease, orchitis/epididymitis, legionella, lyme disease, influenza a, epstein-barr virus, vital-associated hemaphagocytic syndrome, vital encephalitis/aseptic meningitis, and the like.

The present invention also provides a method for modulating or treating at least one malignant or amyloid related disease in a cell, tissue, organ, animal or patient, including, but
5 not limited to, at least one of: leukemia, acute leukemia, acute lymphoblastic leukemia (ALL), acute lymphocytic leukemia, B-cell, T-cell or FAB ALL, acute myeloid leukemia (AML), acute myelogenous leukemia, chronic myelocytic leukemia (CML), chronic lymphocytic leukemia (CLL), hairy cell leukemia, myelodysplastic syndrome (MDS), a lymphoma, Hodgkin's disease, a malignant lymphoma, non-hodgkin's lymphoma, Burkitt's lymphoma,
10 multiple myeloma, Kaposi's sarcoma, colorectal carcinoma, pancreatic carcinoma, nasopharyngeal carcinoma, malignant histiocytosis, paraneoplastic syndrome/hypercalcemia of malignancy, solid tumors, bladder cancer, breast cancer, colorectal cancer, endometrial cancer, head cancer, neck cancer, hereditary nonpolyposis cancer, Hodgkin's lymphoma, liver cancer, lung cancer, non-small cell lung cancer, ovarian cancer, pancreatic cancer, prostate cancer,
15 renal cell carcinoma, testicular cancer, adenocarcinomas, sarcomas, malignant melanoma, hemangioma, metastatic disease, cancer related bone resorption, cancer related bone pain, and the like. Any method of the present invention can comprise administering an effective amount of a composition or pharmaceutical composition comprising at least one anti-amyloid antibody to a cell, tissue, organ, animal or patient in need of such modulation, treatment or
20 therapy. Such a method can optionally further comprise co-administration or combination therapy for treating such diseases or disorders, wherein the administering of said at least one anti-amyloid antibody, specified portion or variant thereof, further comprises administering, before concurrently, and/or after, at least one selected from at least one TNF antagonist (e.g., but not limited to a TNF chemical or protein antagonist, TNF monoclonal or polyclonal
25 antibody or fragment, a soluble TNF receptor (e.g., p55, p70 or p85) or fragment, fusion polypeptides thereof, or a small molecule TNF antagonist, e.g., TNF binding protein I or II (TBP-I or TBP-II), nerelimonmab, infliximab, etanercept, CDP-571, CDP-870, afelimomab, lenercept, and the like), an antirheumatic (e.g., methotrexate, auranofin, aurothioglucose, azathioprine, etanercept, gold sodium thiomalate, hydroxychloroquine sulfate, leflunomide,
30 sulfasalazine), a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial (e.g., aminoglycoside, an antifungal, an antiparasitic, an antiviral, a carbapenem, cephalosporin, a fluorquinolone, a macrolide, a penicillin, a sulfonamide, a tetracycline, another antimicrobial), an antipsoriatic, a corticosteroid, an anabolic steroid, a diabetes related
35 agent, a mineral, a nutritional, a thyroid agent, a vitamin, a calcium related hormone, an

antidiarrheal, an antitussive, an antiemetic, an antiulcer, a laxative, an anticoagulant, an erythropoietin (e.g., epoetin alpha), a filgrastim (e.g., G-CSF, Neupogen), a sargramostim (GM-CSF, Leukine), an immunization, an immunoglobulin, an immunosuppressive (e.g., basiliximab, cyclosporine, daclizumab), a growth hormone, a hormone replacement drug, an estrogen receptor modulator, a mydriatic, a cycloplegic, an alkylating agent, an antimetabolite, a mitotic inhibitor, a radiopharmaceutical, an antidepressant, antimanic agent, an antipsychotic, an anxiolytic, a hypnotic, a sympathomimetic, a stimulant, donepezil, tacrine, an asthma medication, a beta agonist, an inhaled steroid, a leukotriene inhibitor, a methylxanthine, a cromolyn, an epinephrine or analog, dornase alpha (Pulmozyme), a cytokine or a cytokine antagonist. Suitable dosages are well known in the art. See, e.g., Wells et al., eds., Pharmacotherapy Handbook, 2nd Edition, Appleton and Lange, Stamford, CT (2000); PDR Pharmacopoeia, Tarascon Pocket Pharmacopoeia 2000, Deluxe Edition, Tarascon Publishing, Loma Linda, CA (2000); Nursing 2001 Handbook of Drugs, 21st edition, Springhouse Corp., Springhouse, PA, 2001; Health Professional's Drug Guide 2001, ed., Shannon, Wilson, Stang, Prentice-Hall, Inc, Upper Saddle River, NJ. each of which references are entirely incorporated herein by reference.

TNF antagonists suitable for compositions, combination therapy, co-administration, devices and/or methods of the present invention (further comprising at least one anti body, specified portion and variant thereof, of the present invention), include, but are not limited to, anti-TNF antibodies, antigen-binding fragments thereof, and receptor molecules which bind specifically to TNF; compounds which prevent and/or inhibit TNF synthesis, TNF release or its action on target cells, such as thalidomide, tenidap, phosphodiesterase inhibitors (e.g., pentoxifylline and rolipram), A2b adenosine receptor agonists and A2b adenosine receptor enhancers; compounds which prevent and/or inhibit TNF receptor signalling, such as mitogen activated protein (MAP) kinase inhibitors; compounds which block and/or inhibit membrane TNF cleavage, such as metalloproteinase inhibitors; compounds which block and/or inhibit TNF activity, such as angiotensin converting enzyme (ACE) inhibitors (e.g., captopril); and compounds which block and/or inhibit TNF production and/or synthesis, such as MAP kinase inhibitors.

As used herein, a "tumor necrosis factor antibody," "TNF antibody," "TNF α antibody," or fragment and the like decreases, blocks, inhibits, abrogates or interferes with TNF α activity *in vitro*, *in situ* and/or preferably *in vivo*. For example, a suitable TNF human antibody of the present invention can bind TNF α and includes anti-TNF antibodies, antigen-binding fragments thereof, and specified mutants or domains thereof that bind specifically to TNF α . A suitable TNF antibody or fragment can also decrease block, abrogate, interfere,

prevent and/or inhibit TNF RNA, DNA or protein synthesis, TNF release, TNF receptor signaling, membrane TNF cleavage, TNF activity, TNF production and/or synthesis.

Chimeric antibody cA2 consists of the antigen binding variable region of the high-affinity neutralizing mouse anti-human TNF α IgG1 antibody, designated A2, and the constant regions of a human IgG1, kappa immunoglobulin. The human IgG1 Fc region improves
5 allogeneic antibody effector function, increases the circulating serum half-life and decreases the immunogenicity of the antibody. The avidity and epitope specificity of the chimeric antibody cA2 is derived from the variable region of the murine antibody A2. In a particular embodiment, a preferred source for nucleic acids encoding the variable region of the murine
10 antibody A2 is the A2 hybridoma cell line.

Chimeric A2 (cA2) neutralizes the cytotoxic effect of both natural and recombinant human TNF α in a dose dependent manner. From binding assays of chimeric antibody cA2 and recombinant human TNF α , the affinity constant of chimeric antibody cA2 was calculated to be $1.04 \times 10^{10} \text{ M}^{-1}$. Preferred methods for determining monoclonal antibody specificity and affinity
15 by competitive inhibition can be found in Harlow, *et al.*, *antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1988; Colligan *et al.*, eds., *Current Protocols in Immunology*, Greene Publishing Assoc. and Wiley Interscience, New York, (1992-2000); Kozbor *et al.*, *Immunol. Today*, 4:72-79 (1983); Ausubel *et al.*, eds. *Current Protocols in Molecular Biology*, Wiley Interscience, New York (1987-2000); and
20 Muller, *Meth. Enzymol.*, 92:589-601 (1983), which references are entirely incorporated herein by reference.

In a particular embodiment, murine monoclonal antibody A2 is produced by a cell line designated c134A. Chimeric antibody cA2 is produced by a cell line designated c168A.

Additional examples of monoclonal anti-TNF antibodies that can be used in the
25 present invention are described in the art (see, e.g., U.S. Patent No. 5,231,024; Möller, A. *et al.*, *Cytokine* 2(3):162-169 (1990); U.S. Application No. 07/943,852 (filed September 11, 1992); Rathjen *et al.*, International Publication No. WO 91/02078 (published February 21, 1991); Rubin *et al.*, EPO Patent Publication No. 0 218 868 (published April 22, 1987); Yone *et al.*, EPO Patent Publication No. 0 288 088 (October 26, 1988); Liang, *et al.*, *Biochem. Biophys. Res. Comm.* 137:847-854 (1986); Meager, *et al.*, *Hybridoma* 6:305-311 (1987); Fendly *et al.*,
30 *Hybridoma* 6:359-369 (1987); Bringman, *et al.*, *Hybridoma* 6:489-507 (1987); and Hirai, *et al.*, *J. Immunol. Meth.* 96:57-62 (1987), which references are entirely incorporated herein by reference).

35 **TNF Receptor Molecules**

Preferred TNF receptor molecules useful in the present invention are those that bind TNF α with high affinity (see, e.g., Feldmann *et al.*, International Publication No. WO 92/07076 (published April 30, 1992); Schall *et al.*, *Cell* 61:361-370 (1990); and Loetscher *et al.*, *Cell* 61:351-359 (1990), which references are entirely incorporated herein by reference) and optionally possess low immunogenicity. In particular, the 55 kDa (p55 TNF-R) and the 75 kDa (p75 TNF-R) TNF cell surface receptors are useful in the present invention. Truncated forms of these receptors, comprising the extracellular domains (ECD) of the receptors or functional portions thereof (see, e.g., Corcoran *et al.*, *Eur. J. Biochem.* 223:831-840 (1994)), are also useful in the present invention. Truncated forms of the TNF receptors, comprising ECD, have been detected in urine and serum as 30 kDa and 40 kDa TNF α inhibitory binding proteins (Engelmann, H. *et al.*, *J. Biol. Chem.* 265:1531-1536 (1990)). TNF receptor multimeric molecules and TNF immunoreceptor fusion molecules, and derivatives and fragments or portions thereof, are additional examples of TNF receptor molecules which are useful in the methods and compositions of the present invention. The TNF receptor molecules which can be used in the invention are characterized by their ability to treat patients for extended periods with good to excellent alleviation of symptoms and low toxicity. Low immunogenicity and/or high affinity, as well as other undefined properties, can contribute to the therapeutic results achieved.

TNF receptor multimeric molecules useful in the present invention comprise all or a functional portion of the ECD of two or more TNF receptors linked via one or more polypeptide linkers or other nonpeptide linkers, such as polyethylene glycol (PEG). The multimeric molecules can further comprise a signal peptide of a secreted protein to direct expression of the multimeric molecule. These multimeric molecules and methods for their production have been described in U.S. Application No. 08/437,533 (filed May 9, 1995), the content of which is entirely incorporated herein by reference.

TNF immunoreceptor fusion molecules useful in the methods and compositions of the present invention comprise at least one portion of one or more immunoglobulin molecules and all or a functional portion of one or more TNF receptors. These immunoreceptor fusion molecules can be assembled as monomers, or hetero- or homo-multimers. The immunoreceptor fusion molecules can also be monovalent or multivalent. An example of such a TNF immunoreceptor fusion molecule is TNF receptor/IgG fusion protein. TNF immunoreceptor fusion molecules and methods for their production have been described in the art (Lesslauer *et al.*, *Eur. J. Immunol.* 21:2883-2886 (1991); Ashkenazi *et al.*, *Proc. Natl. Acad. Sci. USA* 88:10535-10539 (1991); Peppel *et al.*, *J. Exp. Med.* 174:1483-1489 (1991); Kolls *et al.*, *Proc. Natl. Acad. Sci. USA* 91:215-219 (1994); Butler *et al.*, *Cytokine* 6(6):616-

623 (1994); Baker *et al.*, *Eur. J. Immunol.* 24:2040-2048 (1994); Beutler *et al.*, U.S. Patent No. 5,447,851; and U.S. Application No. 08/442,133 (filed May 16, 1995), each of which references are entirely incorporated herein by reference). Methods for producing immunoreceptor fusion molecules can also be found in Capon *et al.*, U.S. Patent No. 5,116,964; Capon *et al.*, U.S. Patent No. 5,225,538; and Capon *et al.*, *Nature* 337:525-531 (1989), which references are entirely incorporated herein by reference.

A functional equivalent, derivative, fragment or region of TNF receptor molecule refers to the portion of the TNF receptor molecule, or the portion of the TNF receptor molecule sequence which encodes TNF receptor molecule, that is of sufficient size and sequences to functionally resemble TNF receptor molecules that can be used in the present invention (e.g., bind TNF α with high affinity and possess low immunogenicity). A functional equivalent of TNF receptor molecule also includes modified TNF receptor molecules that functionally resemble TNF receptor molecules that can be used in the present invention (e.g., bind TNF α with high affinity and possess low immunogenicity). For example, a functional equivalent of TNF receptor molecule can contain a "SILENT" codon or one or more amino acid substitutions, deletions or additions (e.g., substitution of one acidic amino acid for another acidic amino acid; or substitution of one codon encoding the same or different hydrophobic amino acid for another codon encoding a hydrophobic amino acid). See Ausubel *et al.*, *Current Protocols in Molecular Biology*, Greene Publishing Assoc. and Wiley-Interscience, New York (1987-2000).

Cytokines include any known cytokine. See, e.g., CopewithCytokines.com. Cytokine antagonists include, but are not limited to, any antibody, fragment or mimetic, any soluble receptor, fragment or mimetic, any small molecule antagonist, or any combination thereof.

Therapeutic Treatments. Any method of the present invention can comprise a method for treating an amyloid mediated disorder, comprising administering an effective amount of a composition or pharmaceutical composition comprising at least one anti-amyloid antibody to a cell, tissue, organ, animal or patient in need of such modulation, treatment or therapy.

Such a method can optionally further comprise co-administration or combination therapy for treating such diseases or disorders, wherein the administering of said at least one anti-amyloid antibody, specified portion or variant thereof, further comprises administering, before concurrently, and/or after, at least one selected from an anti-infective drug, a cardiovascular (CV) system drug, a central nervous system (CNS) drug, an autonomic nervous system (ANS) drug, a respiratory tract drug, a gastrointestinal (GI) tract drug, a hormonal drug, a drug for fluid or electrolyte balance, a hematologic drug, an antineoplastic, an

immunomodulation drug, an ophthalmic, otic or nasal drug, a topical drug, a nutritional drug or the like, at least one TNF antagonist (e.g., but not limited to a TNF antibody or fragment, a soluble TNF receptor or fragment, fusion proteins thereof, or a small molecule TNF antagonist), an antirheumatic (e.g., methotrexate, auranofin, aurothioglucose, azathioprine, etanercept, gold sodium thiomalate, hydroxychloroquine sulfate, leflunomide, sulfasalazine), a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial (e.g., aminoglycoside, an antifungal, an antiparasitic, an antiviral, a carbapenem, cephalosporin, a fluroquinolone, a macrolide, a penicillin, a sulfonamide, a tetracycline, another antimicrobial), an antipsoriatic, a corticosteroid, an anabolic steroid, a diabetes related agent, a mineral, a nutritional, a thyroid agent, a vitamin, a calcium related hormone, an antidiarrheal, an antitussive, an antiemetic, an antiulcer, a laxative, an anticoagulant, an erythropoietin (e.g., epoetin alpha), a filgrastim (e.g., G-CSF, Neupogen), a sargramostim (GM-CSF, Leukine), an immunization, an immunoglobulin, an immunosuppressive (e.g., basiliximab, cyclosporine, daclizumab), a growth hormone, a hormone replacement drug, an estrogen receptor modulator, a mydriatic, a cycloplegic, an alkylating agent, an antimetabolite, a mitotic inhibitor, a radiopharmaceutical, an antidepressant, an antimanic agent, an antipsychotic, an anxiolytic, a hypnotic, a sympathomimetic, a stimulant, donepezil, tacrine, an asthma medication, a beta agonist, an inhaled steroid, a leukotriene inhibitor, a methylxanthine, a cromolyn, an epinephrine or analog, dornase alpha (Pulmozyme), a cytokine or a cytokine antagonist. Such drugs are well known in the art, including formulations, indications, dosing and administration for each presented herein (see., e.g., Nursing 2001 Handbook of Drugs, 21st edition, Springhouse Corp., Springhouse, PA, 2001; Health Professional's Drug Guide 2001, ed., Shannon, Wilson, Stang, Prentice-Hall, Inc, Upper Saddle River, NJ; Pharmacotherapy Handbook, Wells et al., ed., Appleton & Lange, Stamford, CT, each entirely incorporated herein by reference).

Typically, treatment of pathologic conditions is effected by administering an effective amount or dosage of at least one anti-amyloid antibody composition that total, on average, a range from at least about 0.01 to 500 milligrams of at least one anti-amyloid antibody per kilogram of patient per dose, and preferably from at least about 0.1 to 100 milligrams antibody /kilogram of patient per single or multiple administration, depending upon the specific activity of contained in the composition. Alternatively, the effective serum concentration can comprise 0.1-5000 µg/ml serum concentration per single or multiple administration. Suitable dosages are known to medical practitioners and will, of course, depend upon the particular disease state, specific activity of the composition being administered, and the particular patient undergoing treatment. In some

instances, to achieve the desired therapeutic amount, it can be necessary to provide for repeated administration, *i.e.*, repeated individual administrations of a particular monitored or metered dose, where the individual administrations are repeated until the desired daily dose or effect is achieved.

5 Preferred doses can optionally include 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 and/or 100-500 mg/kg/administration, or any
10 range, value or fraction thereof, or to achieve a serum concentration of 0.1, 0.5, 0.9, 1.0, 1.1, 1.2, 1.5, 1.9, 2.0, 2.5, 2.9, 3.0, 3.5, 3.9, 4.0, 4.5, 4.9, 5.0, 5.5, 5.9, 6.0, 6.5, 6.9, 7.0, 7.5, 7.9, 8.0, 8.5, 8.9, 9.0, 9.5, 9.9, 10, 10.5, 10.9, 11, 11.5, 11.9, 12, 12.5, 12.9, 13.0, 13.5, 13.9, 14.0, 14.5, 4.9, 5.0, 5.5, 5.9, 6.0, 6.5, 6.9, 7.0, 7.5, 7.9, 8.0, 8.5, 8.9, 9.0, 9.5, 9.9, 10, 10.5, 10.9, 11, 11.5, 11.9, 12, 12.5, 12.9, 13.0, 13.5, 13.9, 14, 14.5, 15, 15.5, 15.9, 16, 16.5, 16.9, 17, 17.5, 17.9, 18, 18.5, 18.9, 19, 19.5, 19.9, 20, 20.5, 20.9, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 96, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, and/or 5000 µg/ml serum concentration per single or multiple
15 administration, or any range, value or fraction thereof.

Alternatively, the dosage administered can vary depending upon known factors, such
20 as the pharmacodynamic characteristics of the particular agent, and its mode and route of administration; age, health, and weight of the recipient; nature and extent of symptoms, kind of concurrent treatment, frequency of treatment, and the effect desired. Usually a dosage of active ingredient can be about 0.1 to 100 milligrams per kilogram of body weight. Ordinarily 0.1 to 50, and preferably 0.1 to 10 milligrams per kilogram per administration or in sustained
25 release form is effective to obtain desired results.

As a non-limiting example, treatment of humans or animals can be provided as a one-time or periodic dosage of at least one antibody of the present invention 0.1 to 100 mg/kg, such as 0.5, 0.9, 1.0, 1.1, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, 45, 50, 60, 70, 80, 90 or 100 mg/kg, per day, on at least one
30 of day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40, or alternatively or additionally, at least one of week 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, or 52, or alternatively or additionally, at least one of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 years, or any combination thereof, using single, infusion or
35

repeated doses.

Dosage forms (composition) suitable for internal administration generally contain from about 0.001 milligram to about 500 milligrams of active ingredient per unit or container. In these pharmaceutical compositions the active ingredient will ordinarily be present in an amount of about 0.5-99.999% by weight based on the total weight of the composition.

For parenteral administration, the antibody can be formulated as a solution, suspension, emulsion, particle, powder, or lyophilized powder in association, or separately provided, with a pharmaceutically acceptable parenteral vehicle. Examples of such vehicles are water, saline, Ringer's solution, dextrose solution, and 1-10% human serum albumin. Liposomes and nonaqueous vehicles such as fixed oils can also be used. The vehicle or lyophilized powder can contain additives that maintain isotonicity (e.g., sodium chloride, mannitol) and chemical stability (e.g., buffers and preservatives). The formulation is sterilized by known or suitable techniques.

Suitable pharmaceutical carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, A. Osol, a standard reference text in this field.

Alternative Administration

Many known and developed modes of can be used according to the present invention for administering pharmaceutically effective amounts of at least one anti-amyloid antibody according to the present invention. While pulmonary administration is used in the following description, other modes of administration can be used according to the present invention with suitable results.

Amyloid antibodies of the present invention can be delivered in a carrier, as a solution, emulsion, colloid, or suspension, or as a dry powder, using any of a variety of devices and methods suitable for administration by inhalation or other modes described here within or known in the art.

Parenteral Formulations and Administration

Formulations for parenteral administration can contain as common excipients sterile water or saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes and the like. Aqueous or oily suspensions for injection can be prepared by using an appropriate emulsifier or humidifier and a suspending agent, according to known methods. Agents for injection can be a non-toxic, non-orally administrable diluting agent such as aqueous solution or a sterile injectable solution or suspension in a solvent. As the usable vehicle or solvent, water, Ringer's solution, isotonic saline, etc. are allowed; as an ordinary solvent, or suspending solvent, sterile involatile oil can be used. For these purposes, any kind of involatile oil and fatty acid can be used, including natural or synthetic or

semisynthetic fatty oils or fatty acids; natural or synthetic or semisynthetic mono- or di- or tri-glycerides. Parental administration is known in the art and includes, but is not limited to, conventional means of injections, a gas pressured needle-less injection device as described in U.S. Pat. No. 5,851,198, and a laser perforator device as described in U.S. Pat. No. 5,839,446
5 entirely incorporated herein by reference.

Alternative Delivery

The invention further relates to the administration of at least one anti-amyloid antibody by parenteral, subcutaneous, intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelial, intracelebellar,
10 intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, intralesional, bolus, vaginal, rectal, buccal, sublingual, intranasal, or transdermal means. At least one anti-amyloid antibody composition can be prepared for use for
15 parenteral (subcutaneous, intramuscular or intravenous) or any other administration particularly in the form of liquid solutions or suspensions; for use in vaginal or rectal administration particularly in semisolid forms such as, but not limited to, creams and suppositories; for buccal, or sublingual administration such as, but not limited to, in the form of tablets or capsules; or intranasally such as, but not limited to, the form of powders, nasal
20 drops or aerosols or certain agents; or transdermally such as not limited to a gel, ointment, lotion, suspension or patch delivery system with chemical enhancers such as dimethyl sulfoxide to either modify the skin structure or to increase the drug concentration in the transdermal patch (Junginger, et al. In "Drug Permeation Enhancement"; Hsieh, D. S., Eds., pp. 59-90 (Marcel Dekker, Inc. New York 1994, entirely incorporated herein by reference), or with
25 oxidizing agents that enable the application of formulations containing proteins and peptides onto the skin (WO 98/53847), or applications of electric fields to create transient transport pathways such as electroporation, or to increase the mobility of charged drugs through the skin such as iontophoresis, or application of ultrasound such as sonophoresis (U.S. Pat. Nos. 4,309,989 and 4,767,402) (the above publications and patents being entirely incorporated
30 herein by reference).

Pulmonary/Nasal Administration

For pulmonary administration, preferably at least one anti-amyloid antibody composition is delivered in a particle size effective for reaching the lower airways of the lung or sinuses. According to the invention, at least one anti-amyloid antibody can be delivered by
35 any of a variety of inhalation or nasal devices known in the art for administration of a

therapeutic agent by inhalation. These devices capable of depositing aerosolized formulations in the sinus cavity or alveoli of a patient include metered dose inhalers, nebulizers, dry powder generators, sprayers, and the like. Other devices suitable for directing the pulmonary or nasal administration of antibodies are also known in the art. All such devices can use of

5 formulations suitable for the administration for the dispensing of antibody in an aerosol. Such aerosols can be comprised of either solutions (both aqueous and non aqueous) or solid particles. Metered dose inhalers like the Ventolin[®] metered dose inhaler, typically use a propellant gas and require actuation during inspiration (See, e.g., WO 94/16970, WO 98/35888). Dry powder inhalers like Turbuhaler[™] (Astra), Rotahaler[®] (Glaxo), Diskus[®]

10 (Glaxo), Spiros[™] inhaler (Dura), devices marketed by Inhale Therapeutics, and the Spinhaler[®] powder inhaler (Fisons), use breath-actuation of a mixed powder (US 4668218 Astra, EP 237507 Astra, WO 97/25086 Glaxo, WO 94/08552 Dura, US 5458135 Inhale, WO 94/06498 Fisons, entirely incorporated herein by reference). Nebulizers like AERx[™] Aradigm, the Ultravent[®] nebulizer (Mallinckrodt), and the Acorn II[®] nebulizer (Marquest Medical Products)

15 (US 5404871 Aradigm, WO 97/22376), the above references entirely incorporated herein by reference, produce aerosols from solutions, while metered dose inhalers, dry powder inhalers, etc. generate small particle aerosols. These specific examples of commercially available inhalation devices are intended to be a representative of specific devices suitable for the practice of this invention, and are not intended as limiting the scope of the invention.

20 Preferably, a composition comprising at least one anti-amyloid antibody is delivered by a dry powder inhaler or a sprayer. There are a several desirable features of an inhalation device for administering at least one antibody of the present invention. For example, delivery by the inhalation device is advantageously reliable, reproducible, and accurate. The inhalation device can optionally deliver small dry particles, e.g. less than about 10 μm , preferably about 1-5 μm ,

25 for good respirability.

Administration of amyloid antibody Compositions as a Spray

A spray including amyloid antibody composition can be produced by forcing a suspension or solution of at least one anti-amyloid antibody through a nozzle under pressure. The nozzle size and configuration, the applied pressure, and the liquid feed rate can be chosen

30 to achieve the desired output and particle size. An electrospray can be produced, for example, by an electric field in connection with a capillary or nozzle feed. Advantageously, particles of at least one anti-amyloid antibody composition delivered by a sprayer have a particle size less than about 10 μm , preferably in the range of about 1 μm to about 5 μm , and most preferably about 2 μm to about 3 μm .

35 Formulations of at least one anti-amyloid antibody composition suitable for use with a

sprayer typically include antibody composition in an aqueous solution at a concentration of about 0.1 mg to about 100 mg of at least one anti-amyloid antibody composition per ml of solution or mg/gm, or any range or value therein, e.g., but not limited to, .1, .2, .3, .4, .5, .6, .7, .8, .9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, 45, 50, 60, 70, 80, 90 or 100 mg/ml or mg/gm. The formulation can include agents such as an excipient, a buffer, an isotonicity agent, a preservative, a surfactant, and, preferably, zinc. The formulation can also include an excipient or agent for stabilization of the antibody composition, such as a buffer, a reducing agent, a bulk protein, or a carbohydrate. Bulk proteins useful in formulating antibody compositions include albumin, protamine, or the like. Typical carbohydrates useful in formulating antibody compositions include sucrose, mannitol, lactose, trehalose, glucose, or the like. The antibody composition formulation can also include a surfactant, which can reduce or prevent surface-induced aggregation of the antibody composition caused by atomization of the solution in forming an aerosol. Various conventional surfactants can be employed, such as polyoxyethylene fatty acid esters and alcohols, and polyoxyethylene sorbitol fatty acid esters. Amounts will generally range between 0.001 and 14% by weight of the formulation. Especially preferred surfactants for purposes of this invention are polyoxyethylene sorbitan monooleate, polysorbate 80, polysorbate 20, or the like. Additional agents known in the art for formulation of a protein such as amyloid antibodies, or specified portions or variants, can also be included in the formulation.

Administration of amyloid antibody compositions by a Nebulizer

Antibody composition can be administered by a nebulizer, such as jet nebulizer or an ultrasonic nebulizer. Typically, in a jet nebulizer, a compressed air source is used to create a high-velocity air jet through an orifice. As the gas expands beyond the nozzle, a low-pressure region is created, which draws a solution of antibody composition through a capillary tube connected to a liquid reservoir. The liquid stream from the capillary tube is sheared into unstable filaments and droplets as it exits the tube, creating the aerosol. A range of configurations, flow rates, and baffle types can be employed to achieve the desired performance characteristics from a given jet nebulizer. In an ultrasonic nebulizer, high-frequency electrical energy is used to create vibrational, mechanical energy, typically employing a piezoelectric transducer. This energy is transmitted to the formulation of antibody composition either directly or through a coupling fluid, creating an aerosol including the antibody composition. Advantageously, particles of antibody composition delivered by a nebulizer have a particle size less than about 10 μm , preferably in the range of about 1 μm to about 5 μm , and most preferably about 2 μm to about 3 μm .

Formulations of at least one anti-amyloid antibody suitable for use with a nebulizer, either jet or ultrasonic, typically include a concentration of about 0.1 mg to about 100 mg of at least one anti-amyloid antibody protein per ml of solution. The formulation can include agents such as an excipient, a buffer, an isotonicity agent, a preservative, a surfactant, and, preferably, zinc. The formulation can also include an excipient or agent for stabilization of the at least one anti-amyloid antibody composition, such as a buffer, a reducing agent, a bulk protein, or a carbohydrate. Bulk proteins useful in formulating at least one anti-amyloid antibody compositions include albumin, protamine, or the like. Typical carbohydrates useful in formulating at least one anti-amyloid antibody include sucrose, mannitol, lactose, trehalose, glucose, or the like. The at least one anti-amyloid antibody formulation can also include a surfactant, which can reduce or prevent surface-induced aggregation of the at least one anti-amyloid antibody caused by atomization of the solution in forming an aerosol. Various conventional surfactants can be employed, such as polyoxyethylene fatty acid esters and alcohols, and polyoxyethylene sorbital fatty acid esters. Amounts will generally range between 0.001 and 4% by weight of the formulation. Especially preferred surfactants for purposes of this invention are polyoxyethylene sorbitan mono-oleate, polysorbate 80, polysorbate 20, or the like. Additional agents known in the art for formulation of a protein such as antibody protein can also be included in the formulation.

Administration of amyloid antibody compositions By A Metered Dose Inhaler

In a metered dose inhaler (MDI), a propellant, at least one anti-amyloid antibody, and any excipients or other additives are contained in a canister as a mixture including a liquefied compressed gas. Actuation of the metering valve releases the mixture as an aerosol, preferably containing particles in the size range of less than about 10 μm , preferably about 1 μm to about 5 μm , and most preferably about 2 μm to about 3 μm . The desired aerosol particle size can be obtained by employing a formulation of antibody composition produced by various methods known to those of skill in the art, including jet-milling, spray drying, critical point condensation, or the like. Preferred metered dose inhalers include those manufactured by 3M or Glaxo and employing a hydrofluorocarbon propellant.

Formulations of at least one anti-amyloid antibody for use with a metered-dose inhaler device will generally include a finely divided powder containing at least one anti-amyloid antibody as a suspension in a non-aqueous medium, for example, suspended in a propellant with the aid of a surfactant. The propellant can be any conventional material employed for this purpose, such as chlorofluorocarbon, a hydrochlorofluorocarbon, a hydrofluorocarbon, or a hydrocarbon, including trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethanol and 1,1,1,2-tetrafluoroethane, HFA-134a (hydrofluoroalkane-134a),

HFA-227 (hydrofluoroalkane-227), or the like. Preferably the propellant is a hydrofluorocarbon. The surfactant can be chosen to stabilize the at least one anti-amyloid antibody as a suspension in the propellant, to protect the active agent against chemical degradation, and the like. Suitable surfactants include sorbitan trioleate, soya lecithin, oleic acid, or the like. In some cases solution aerosols are preferred using solvents such as ethanol. Additional agents known in the art for formulation of a protein such as protein can also be included in the formulation.

One of ordinary skill in the art will recognize that the methods of the current invention can be achieved by pulmonary administration of at least one anti-amyloid antibody compositions via devices not described herein.

Oral Formulations and Administration

Formulations for oral rely on the co-administration of adjuvants (e.g., resorcinols and nonionic surfactants such as polyoxyethylene oleyl ether and n-hexadecylpolyethylene ether) to increase artificially the permeability of the intestinal walls, as well as the co-administration of enzymatic inhibitors (e.g., pancreatic trypsin inhibitors, diisopropylfluorophosphate (DFF) and trasylol) to inhibit enzymatic degradation. Formulations for delivery of hydrophilic agents including proteins and antibodies and a combination of at least two surfactants intended for oral, buccal, mucosal, nasal, pulmonary, vaginal transmembrane, or rectal administration are taught in U.S. 6,309,663. The active constituent compound of the solid-type dosage form for oral administration can be mixed with at least one additive, including sucrose, lactose, cellulose, mannitol, trehalose, raffinose, maltitol, dextran, starches, agar, arginates, chitins, chitosans, pectins, gum tragacanth, gum arabic, gelatin, collagen, casein, albumin, synthetic or semisynthetic polymer, and glyceride. These dosage forms can also contain other type(s) of additives, e.g., inactive diluting agent, lubricant such as magnesium stearate, paraben, preserving agent such as sorbic acid, ascorbic acid, alpha-tocopherol, antioxidant such as cysteine, disintegrator, binder, thickener, buffering agent, sweetening agent, flavoring agent, perfuming agent, etc.

Tablets and pills can be further processed into enteric-coated preparations. The liquid preparations for oral administration include emulsion, syrup, elixir, suspension and solution preparations allowable for medical use. These preparations can contain inactive diluting agents ordinarily used in said field, e.g., water. Liposomes have also been described as drug delivery systems for insulin and heparin (U.S. Pat. No. 4,239,754). More recently, microspheres of artificial polymers of mixed amino acids (proteinoids) have been used to deliver pharmaceuticals (U.S. Pat. No. 4,925,673). Furthermore, carrier compounds described in U.S. Pat. No. 5,879,681 and U.S. Pat. No. 5,587,753 are used to deliver biologically active agents

orally are known in the art.

Mucosal Formulations and Administration

A formulation for orally administering a bioactive agent encapsulated in one or more biocompatible polymer or copolymer excipients, preferably a biodegradable polymer or copolymer, affording microcapsules which due to the proper size of the resultant microcapsules results in the agent reaching and being taken up by the folliculi lymphatic aggregati, otherwise known as the "Peyer's patch," or "GALT" of the animal without loss of effectiveness due to the agent having passed through the gastrointestinal tract. Similar folliculi lymphatic aggregati can be found in the bronchi tubes (BALT) and the large intestine. The above-described tissues are referred to in general as mucosally associated lymphoreticular tissues (MALT). For absorption through mucosal surfaces, compositions and methods of administering at least one anti-amyloid antibody include an emulsion comprising a plurality of submicron particles, a mucoadhesive macromolecule, a bioactive peptide, and an aqueous continuous phase, which promotes absorption through mucosal surfaces by achieving mucoadhesion of the emulsion particles (U.S. Pat. No. 5,514,670). Mucous surfaces suitable for application of the emulsions of the present invention can include corneal, conjunctival, buccal, sublingual, nasal, vaginal, pulmonary, stomachic, intestinal, and rectal routes of administration. Formulations for vaginal or rectal administration, e.g. suppositories, can contain as excipients, for example, polyalkyleneglycols, vaseline, cocoa butter, and the like. Formulations for intranasal administration can be solid and contain as excipients, for example, lactose or can be aqueous or oily solutions of nasal drops. For buccal administration excipients include sugars, calcium stearate, magnesium stearate, pregelatinized starch, and the like (U.S. Pat. No. 5,849,695).

Transdermal Formulations and Administration

For transdermal administration, the at least one anti-amyloid antibody is encapsulated in a delivery device such as a liposome or polymeric nanoparticles, microparticle, microcapsule, or microspheres (referred to collectively as microparticles unless otherwise stated). A number of suitable devices are known, including microparticles made of synthetic polymers such as polyhydroxy acids such as polylactic acid, polyglycolic acid and copolymers thereof, polyorthoesters, polyanhydrides, and polyphosphazenes, and natural polymers such as collagen, polyamino acids, albumin and other proteins, alginate and other polysaccharides, and combinations thereof (U.S. Pat. No. 5,814,599).

Prolonged Administration and Formulations

It can be sometimes desirable to deliver the compounds of the present invention to the subject over prolonged periods of time, for example, for periods of one week to one year from

a single administration. Various slow release, depot or implant dosage forms can be utilized. For example, a dosage form can contain a pharmaceutically acceptable non-toxic salt of the compounds that has a low degree of solubility in body fluids, for example, (a) an acid addition salt with a polybasic acid such as phosphoric acid, sulfuric acid, citric acid, tartaric acid, tannic acid, pamoic acid, alginic acid, polyglutamic acid, naphthalene mono- or di-sulfonic acids, polygalacturonic acid, and the like; (b) a salt with a polyvalent metal cation such as zinc, calcium, bismuth, barium, magnesium, aluminum, copper, cobalt, nickel, cadmium and the like, or with an organic cation formed from e.g., N,N'-dibenzyl-ethylenediamine or ethylenediamine; or (c) combinations of (a) and (b) e.g. a zinc tannate salt. Additionally, the compounds of the present invention or, preferably, a relatively insoluble salt such as those just described, can be formulated in a gel, for example, an aluminum monostearate gel with, e.g. sesame oil, suitable for injection. Particularly preferred salts are zinc salts, zinc tannate salts, pamoate salts, and the like. Another type of slow release depot formulation for injection would contain the compound or salt dispersed for encapsulated in a slow degrading, non-toxic, non-antigenic polymer such as a polylactic acid/polyglycolic acid polymer for example as described in U.S. Pat. No. 3,773,919. The compounds or, preferably, relatively insoluble salts such as those described above can also be formulated in cholesterol matrix silastic pellets, particularly for use in animals. Additional slow release, depot or implant formulations, e.g. gas or liquid liposomes are known in the literature (U.S. Pat. No. 5,770,222 and "Sustained and Controlled Release Drug Delivery Systems", J. R. Robinson ed., Marcel Dekker, Inc., N.Y., 1978).

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

Example 1: Cloning and Expression of amyloid antibody in Mammalian Cells

A typical mammalian expression vector contains at least one promoter element, which mediates the initiation of transcription of the antibody coding sequences, encoding heavy and light chain variable regions adjacent to coding sequences of known constant regions, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription can be achieved with the early and late promoters from SV40, the long terminal repeats (LTRS) from Retroviruses, e.g., RSV, HTLV, HIV and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter). Suitable expression vectors for use

in practicing the present invention include, for example, vectors such as pIRES1neo, pRetro-Off, pRetro-On, PLXSN, or pLNCX (Clontech Labs, Palo Alto, CA), pcDNA3.1 (+/-), pcDNA/Zeo (+/-) or pcDNA3.1/Hygro (+/-) (Invitrogen), PSVL and PMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146) and pBC12MI (ATCC 5 67109). Mammalian host cells that could be used include human Hela 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV 1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, the gene can be expressed in stable cell lines that contain the gene integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, 10 neomycin, or hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the encoded antibody. The DHFR (dihydrofolate reductase) marker is useful to develop cell lines that carry several hundred or even several thousand copies of the gene of interest. Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy, et al., Biochem. J. 227:277- 15 279 (1991); Bebbington, et al., Bio/Technology 10:169-175 (1992)). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of antibodies.

The expression vectors pC1 and pC4 contain the strong promoter (LTR) of the Rous 20 Sarcoma Virus (Cullen, et al., Molec. Cell. Biol. 5:438-447 (1985)) plus a fragment of the CMV-enhancer (Boshart, et al., Cell 41:521-530 (1985)). Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors contain in addition the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene.

25 **Cloning and Expression in CHO Cells**

The vector pC4 is used for the expression of amyloid antibody. Plasmid pC4 is a derivative of the plasmid pSV2-dhfr (ATCC Accession No. 37146). The plasmid contains the mouse DHFR gene under control of the SV40 early promoter. Chinese hamster ovary- or other cells lacking dihydrofolate activity that are transfected with these plasmids can be selected by 30 growing the cells in a selective medium (e.g., alpha minus MEM, Life Technologies, Gaithersburg, MD) supplemented with the chemotherapeutic agent methotrexate. The amplification of the DHFR genes in cells resistant to methotrexate (MTX) has been well documented (see, e.g., F. W. Alt, et al., J. Biol. Chem. 253:1357-1370 (1978); J. L. Hamlin and C. Ma, Biochem. et Biophys. Acta 1097:107-143 (1990); and M. J. Page and M. A. Sydenham, 35 Biotechnology 9:64-68 (1991)). Cells grown in increasing concentrations of MTX develop

resistance to the drug by overproducing the target enzyme, DHFR, as a result of amplification of the DHFR gene. If a second gene is linked to the DHFR gene, it is usually co-amplified and over-expressed. It is known in the art that this approach can be used to develop cell lines carrying more than 1,000 copies of the amplified gene(s). Subsequently, when the
5 methotrexate is withdrawn, cell lines are obtained that contain the amplified gene integrated into one or more chromosome(s) of the host cell.

Plasmid pC4 contains for expressing the gene of interest the strong promoter of the long terminal repeat (LTR) of the Rous Sarcoma Virus (Cullen, et al., *Molec. Cell. Biol.* 5:438-447 (1985)) plus a fragment isolated from the enhancer of the immediate early gene of human
10 cytomegalovirus (CMV) (Boshart, et al., *Cell* 41:521-530 (1985)). Downstream of the promoter are BamHI, XbaI, and Asp718 restriction enzyme cleavage sites that allow integration of the genes. Behind these cloning sites the plasmid contains the 3' intron and polyadenylation site of the rat preproinsulin gene. Other high efficiency promoters can also be used for the expression, e.g., the human b-actin promoter, the SV40 early or late promoters or
15 the long terminal repeats from other retroviruses, e.g., HIV and HTLV. Clontech's Tet-Off and Tet-On gene expression systems and similar systems can be used to express the amyloid in a regulated way in mammalian cells (M. Gossen, and H. Bujard, *Proc. Natl. Acad. Sci. USA* 89: 5547-5551 (1992)). For the polyadenylation of the mRNA other signals, e.g., from the human growth hormone or globin genes can be used as well. Stable cell lines carrying a gene
20 of interest integrated into the chromosomes can also be selected upon co-transfection with a selectable marker such as gpt, G418 or hygromycin. It is advantageous to use more than one selectable marker in the beginning, e.g., G418 plus methotrexate.

The plasmid pC4 is digested with restriction enzymes and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The vector is then isolated from a
25 1% agarose gel.

In one set of experiments, the DNA sequence encoding the complete amyloid antibody is used, e.g., as presented in SEQ ID NOS:51 or 52, corresponding to HC and LC variable regions of the amyloid antibody of the present invention as presented in SEQ ID NOS:48 or 49, according to known method steps. Isolated nucleic acid encoding a suitable human constant
30 region (i.e., HC and LC regions) is also used in this construct. In another set of experiments, the DNA sequence as presented in SEQ ID NOS:61 or 62, corresponding to HC and LC variable regions as presented in SEQ ID NOS:59 or 60, is used. The DNA sequence as presented in SEQ ID NOS:71 or 72, corresponding to HC and LC variable regions as presented in SEQ ID NOS:69 or 70, and the DNA sequence as presented in SEQ ID NOS:81 or 82,
35 corresponding to HC and LC variable regions as presented in SEQ ID NOS:79 or 80, are also

used.

The isolated variable and constant region encoding DNA and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC4 using, for instance, restriction enzyme analysis.

Chinese hamster ovary (CHO) cells lacking an active DHFR gene are used for transfection. 5 µg of the expression plasmid pC4 is cotransfected with 0.5 µg of the plasmid pSV2-neo using lipofectin. The plasmid pSV2neo contains a dominant selectable marker, the neo gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 µg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 µg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 mM, 2 mM, 5 mM, 10 mM, 20 mM). The same procedure is repeated until clones are obtained that grow at a concentration of 100 - 200 mM. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reverse phase HPLC analysis.

Binding Kinetics of Human Anti-Human amyloid antibodies

ELISA analysis confirms that purified antibody from these host cells bind amyloid in a concentration-dependent manner. In this case, the avidity of the antibody for its cognate antigen (epitope) is measured. Quantitative binding constants are obtained using BIAcore analysis of the human antibodies and reveals that several of the human monoclonal antibodies are very high affinity with K_D in the range of 1×10^{-9} to 9×10^{-12} .

Conclusions

Human amyloid reactive IgG monoclonal antibodies of the invention are generated. The human anti-amyloid antibodies are further characterized. Several of generated antibodies have affinity constants between 1×10^8 and 9×10^{12} . The high affinities of these fully human monoclonal antibodies make them suitable for therapeutic applications in amyloid-dependent diseases, pathologies or related conditions.

Example 2: Expression and Purification of an amyloid Protein or Antibody in *E. coli*

The bacterial expression vector pQE60 is used for bacterial expression in this example.

(QIAGEN, Inc., Chatsworth, CA). pQE60 encodes ampicillin antibiotic resistance ("Ampr") and contains a bacterial origin of replication ("ori"), an IPTG inducible promoter, a ribosome binding site ("RBS"), six codons encoding histidine residues that allow affinity purification using nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin sold by QIAGEN, Inc., and
5 suitable single restriction enzyme cleavage sites. These elements are arranged such that a DNA fragment encoding a protein or antibody can be inserted in such a way as to produce that protein or antibody with the six His residues (i.e., a "6 X His tag") covalently linked to the carboxyl terminus of that protein or antibody. However, a protein or antibody coding sequence can optionally be inserted such that translation of the six His codons is prevented and,
10 therefore, a protein or antibody is produced with no 6 X His tag.

The nucleic acid sequence encoding the desired portion of an amyloid antibody, e.g., the HC and LC variable region as represented in SEQ ID NOS:48, 49, 59, 60, 69, 70, 79 and 80, the HC CDRs as represented in SEQ ID NOS:42-44, 53-55, 63-65 and 73-75, the LC CDRs as represented in SEQ ID NOS:45-47, 56-58, 66-68, and 76-78, optionally further comprising
15 part or all of the coding sequence for a known human constant region optionally and preferably lacking the hydrophobic leader sequence is amplified from the deposited cDNA clone using PCR oligonucleotide primers (based on the sequences presented, which anneal to the amino terminal encoding DNA sequences of the desired portion of an amyloid protein or antibody and to sequences in the deposited construct 3' to the cDNA coding sequence. Additional
20 nucleotides containing restriction sites to facilitate cloning in the pQE60 vector are added to the 5' and 3' sequences, respectively.

For cloning an amyloid protein or antibody, the 5' and 3' primers have nucleotides corresponding or complementary to a portion of the coding sequence of an amyloid protein or antibody, according to known method steps. One of ordinary skill in the art would appreciate,
25 of course, that the point in a protein or antibody coding sequence where the 5' primer begins can be varied to amplify a desired portion of the complete protein or antibody shorter or longer than the mature form.

The amplified amyloid nucleic acid fragments and the vector pQE60 are digested with appropriate restriction enzymes and the digested DNAs are then ligated together. Insertion of
30 the amyloid DNA into the restricted pQE60 vector places an amyloid protein or antibody coding region including its associated stop codon downstream from the IPTG-inducible promoter and in-frame with an initiating AUG codon. The associated stop codon prevents translation of the six histidine codons downstream of the insertion point.

The ligation mixture is transformed into competent *E. coli* cells using standard
35 procedures such as those described in Sambrook, et al., 1989; Ausubel, 1987-1998. *E. coli* CEN5021 NP

strain MI5/rep4, containing multiple copies of the plasmid pREP4, which expresses the lac repressor and confers kanamycin resistance ("Kan^r"), is used in carrying out the illustrative example described herein. This strain, which is only one of many that are suitable for expressing amyloid protein or antibody, is available commercially from QIAGEN, Inc.

- 5 Transformants are identified by their ability to grow on LB plates in the presence of ampicillin and kanamycin. Plasmid DNA is isolated from resistant colonies and the identity of the cloned DNA confirmed by restriction analysis, PCR and DNA sequencing.

- Clones containing the desired constructs are grown overnight ("O/N") in liquid culture in LB media supplemented with both ampicillin (100 µg/ml) and kanamycin (25 µg/ml). The
- 10 O/N culture is used to inoculate a large culture, at a dilution of approximately 1:25 to 1:250. The cells are grown to an optical density at 600 nm ("OD₆₀₀") of between 0.4 and 0.6. Isopropyl-b-D-thiogalactopyranoside ("IPTG") is then added to a final concentration of 1 mM to induce transcription from the lac repressor sensitive promoter, by inactivating the lacI repressor. Cells subsequently are incubated further for 3 to 4 hours. Cells then are harvested
- 15 by centrifugation.

- The cells are then stirred for 3-4 hours at 4°C in 6M guanidine-HCl, pH8. The cell debris is removed by centrifugation, and the supernatant containing the amyloid is dialyzed against 50 mM Na-acetate buffer pH6, supplemented with 200 mM NaCl. Alternatively, a protein or antibody can be successfully refolded by dialyzing it against 500 mM NaCl, 20%
- 20 glycerol, 25 mM Tris/HCl pH7.4, containing protease inhibitors.

- If insoluble protein is generated, the protein is made soluble according to known method steps. After renaturation the protein or antibody is purified by ion exchange, hydrophobic interaction and size exclusion chromatography. Alternatively, an affinity chromatography step such as an antibody column is used to obtain pure amyloid protein or
- 25 antibody. The purified protein or antibody is stored at 4°C or frozen at -40°C to -120°C.

Example 3: Cloning and Expression of an amyloid Polypeptide in a Baculovirus Expression System

In this illustrative example, the plasmid shuttle vector pA2 GP is used to insert the
5 cloned DNA encoding the antibody (e.g. comprising the variable regions of SEQ ID NOS:51-
52, 61-62, 71-72, or 81-82) into a baculovirus to express an amyloid antibody, using a
baculovirus leader and standard methods as described in Summers, et al., A Manual of
Methods for Baculovirus Vectors and Insect Cell Culture Procedures, Texas Agricultural
Experimental Station Bulletin No. 1555 (1987). This expression vector contains the strong
10 polyhedrin promoter of the Autographa californica nuclear polyhedrosis virus (AcMNPV)
followed by the secretory signal peptide (leader) of the baculovirus gp67 protein or antibody
and convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of
the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of
recombinant virus, the plasmid contains the beta-galactosidase gene from E. coli under control
15 of a weak Drosophila promoter in the same orientation, followed by the polyadenylation signal
of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for
cell-mediated homologous recombination with wild-type viral DNA to generate viable virus
that expresses the cloned polynucleotide.

Other baculovirus vectors are used in place of the vector above, such as pAc373,
20 pVL941 and pAcIM1, as one skilled in the art would readily appreciate, as long as the
construct provides appropriately located signals for transcription, translation, secretion and the
like, including a signal peptide and an in-frame AUG as required. Such vectors are described,
for instance, in Luckow, et al., Virology 170:31-39.

The cDNA sequence encoding the amyloid antibody in the deposited or other clone,
25 lacking the AUG initiation codon and the naturally associated nucleotide binding site, is
amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the
gene. Non-limiting examples include 5' and 3' primers having nucleotides corresponding or
complementary to a portion of the coding sequence of an amyloid protein or antibody, e.g., as
presented in SEQ ID NOS:48-49 for C701, SEQ ID NOS:59-60 for C705, SEQ ID NOS:69-70
30 for C706, or SEQ ID NOS:79-80 for C707, according to known method steps.

The amplified fragment is isolated from a 1% agarose gel using a commercially
available kit (e.g., "GeneClean," BIO 101 Inc., La Jolla, CA). The fragment then is then
digested with the appropriate restriction enzyme and again is purified on a 1% agarose gel.
This fragment is designated herein "F1".

35 The plasmid is digested with the corresponding restriction enzymes and optionally, can
be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the

art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, CA). This vector DNA is designated herein "V1".

Fragment F1 and the dephosphorylated plasmid V1 are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria are identified that contain the plasmid with the human amyloid gene using the PCR method, in which one of the primers that is used to amplify the gene and the second primer is from well within the vector so that only those bacterial colonies containing the amyloid gene fragment will show amplification of the DNA. The sequence of the cloned fragment is confirmed by DNA sequencing. This plasmid is designated herein pBac amyloid .

Five μg of the plasmid pBacamyloid is co-transfected with 1.0 μg of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner, et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). 1 μg of BaculoGold™ virus DNA and 5 μg of the plasmid pBac amyloid are mixed in a sterile well of a microtiter plate containing 50 μl of serum-free Grace's medium (Life Technologies, Inc., Rockville, MD). Afterwards, 10 μl Lipofectin plus 90 μl Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is rocked back and forth to mix the newly added solution. The plate is then incubated for 5 hours at 27°C. After 5 hours the transfection solution is removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. The plate is put back into an incubator and cultivation is continued at 27°C for four days.

After four days the supernatant is collected and a plaque assay is performed, according to known methods. An agarose gel with "Blue Gal" (Life Technologies, Inc., Rockville, MD) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies, Inc., Rockville, MD, page 9-10). After appropriate incubation, blue stained plaques are picked with a micropipettor tip (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 μl of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4°C. The recombinant virus is called V-amyloid.

To verify the expression of the amyloid gene, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus V-amyloid at a multiplicity of infection ("MOI") of about 2. Six hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine
5 (available, e.g., from Life Technologies, Inc., Rockville, MD). If radiolabeled protein or antibodies are desired, 42 hours later, 5 mCi of 35S-methionine and 5 mCi 35S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then they are harvested by centrifugation. The protein or antibodies in the supernatant as well as the intracellular protein or antibodies are analyzed by SDS-PAGE followed by autoradiography (if
10 radiolabeled). Microsequencing of the amino acid sequence of the amino terminus of purified protein or antibody can be used to determine the amino terminal sequence of the mature protein or antibody and thus the cleavage point and length of the secretory signal peptide.

Example 4: Characterization of linear epitopes of anti-human beta amyloid antibodies

1.5 Introduction

Alzheimer's Disease (AD) is a progressive dementia with pathology that can be characterized by hyperphosphorylated tau in neurons and extracellular deposits of beta-amyloid (A β) plaques in the brain. A β plaques form as a result of over-production, or inefficient clearance, of a highly self-aggregating 42 amino acid peptide of amyloid precursor
20 protein (APP). The normal function of APP or the A β ₄₂ peptide is unknown, but it is the A β ₄₂ species that is believed to be related to AD. A β ₄₂ can quickly self-aggregate to form oligomeric structures that progress to fibrils, and eventually plaques. These plaques are a hallmark of AD pathology.

Therapeutic intervention directed towards interrupting the amyloid cascade has been
25 demonstrated in transgenic AD animal models using active vaccination with A β peptide or peripheral administration of A β -specific monoclonal antibodies. The rationale for these approaches relied upon immune system mediated plaque and/or A β clearance. The mechanism of plaque clearance was proposed to occur via direct binding of antibody to plaques within the brain, thus activating microglial cells via Fc receptors to phagocytose the deposited A β .
30 Alternatively, peripherally administered antibodies may bind circulating A β moieties, thus changing the dynamic equilibrium of A β concentrations between the central nervous system and plasma and promoting A β efflux from the brain.

Methods

Peptide synthesis on cellulose membranes

35 Membranes were purchased from Intavis (Bergisch Gladbach, Germany).
CEN5021 NP

Fluorenylmethyloxycarbonyl (Fmoc) amino acids and N-hydroxybenzotriazole (HBOT) were from Novabiochem (Meudon, France). N,N'-diisopropylcarbodiimide (DIC) was from Fluka (Germany). N,N'-dimethylformamide (DMF) and N-methylpyrrolidone-2 (NMP), were obtained from Applied Biosystems. The Rink resin was purchased from Advanced Chem Tech.

5 The peptide synthesis of A β ,

KMDAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVVIATVIVITLVML (SEQ ID NO:50), was performed according to Frank R. (2002) using an Auto Spot Robot ASP 222 (Abimed GmbH, Germany), as previously described (Kramer et al., 1994). The membranes used were derivatized with polyethylene glycol spacer of a length of 8 to 10 ethylene glycol units (Amino-PEG₅₀₀-UC Sheet, loading: 400 nmol/cm²) (Intavis AG, Lot AC112050900).
10 The grid was generated by spotting the C-terminal β -alanine. All peptides were N-acetylated and approximately 20 nmol of peptide per single spot was generated.

Membrane probing and regeneration

After an overnight saturation step in SuperBlock Blocking Buffer (Pierce), protein G
15 purified antibodies were added to the membrane (1 μ g/ml, 2 hrs incubation at room temperature). After washing the membrane, a 1:10,000 dilution of a Horse Radish Peroxidase-conjugated goat anti-mouse IgG (Jackson ImmunoResearch Laboratory Inc.) was incubated for 1 hour at room temperature. Bound antibody was detected by the ECL plus kit (Amersham), which gives a positive indicative of the antibody binding on the spots.

20 Results

A cellulose-bound set of overlapping peptides spanning the primary sequence of human A β was probed with the anti-A β IgG antibodies, C701, C705, C706 and C707. Linear peptide epitopes could be identified for these monoclonal antibodies using a peptide scan (15-mers through 6-mers, with 1 amino acid shift).

25 The Spot membrane results showed that C701 recognized the sequence LMVGGV (Figure 42). C705 recognized the N-terminal sequence, EFRHDS, specifically (Figure 43). C706 recognized the N-terminal sequence, AEFRHD, specifically (Figure 44). C707 recognized the central domain and C-terminal sequence, GLMVGGVIA (Figure 45).

Conclusion

30 In summary, epitope mapping of anti-A β monoclonal antibodies using Spot membranes showed these antibodies recognized linear epitopes. Two mAbs, C705 and C706, are specific for the N-terminal sequence of A β . One mAb, C701 recognizes the C-terminal sequence of A β . Interestingly, the mAb C707 binds the central domain and the C-terminal sequence of A β .

Example 5: Ligand binding of Anti-human beta amyloid antibodies

Methods

BIAcore 3000, CM5 sensor surface, amine coupling kit, HEPES buffered saline (HBS, 10 mM HEPES 150 mM NaCl, pH7.4 with 3 mM EDTA and 0.005% Tween-20) and 10 mM sodium acetate pH 4.5 were purchased from Biacore, Inc. (Piscataway , NJ). Anti-A β monoclonal antibodies (100 ug/mL) were dialyzed against HBS diluted 1:10 with water. Then, the dialyzed mAb solution was diluted 1:10 into 10 mM sodium acetate pH 4.5. The CM5 sensor surface was equilibrated in the BIAcore 3000 with HBS. Each antibody was immobilized onto a flow cell using the immobilization wizard provided in the operating software and the protocols supplied with the amine coupling kit. The wizard was set to immobilize 2500 RU of antibody. Typically 2000 – 3000 RU were actually immobilized.

Results

For each mAb response, data as a function of time were collected (Figure 46). From the sensorgram, report points are taken 10 seconds prior to the injection of the peptide, 10 seconds prior to the completion of the association phase, and 60 seconds into the dissociation phase. This data was then used to determine the binding stoichiometry, and a measure of stability (Fraction of bound peptide remaining on the sensor surface after 60 seconds). These calculations are made possible by assuming that 1 RU = 1 pg of peptide (or protein)/ mm².

Due to the potentially multiple aggregation states of the 1-38 and 1-42 peptides, a quantitative analysis of the binding constants is not possible. However, a qualitative analysis is possible. Figure 47 ranks the mAbs as a function of binding ratio and fraction remaining on the surface after 60 seconds reflecting the stability of the complex. From this analysis C705 and C707 appear to be capable of binding to “monomeric” peptide, where as C701 and C706 requires the peptide to be aggregated to some extent before they can bind.

Example 5: Oligomer neutralization of Anti-human beta amyloid antibodies

Methods

A β ₁₋₄₂ oligomer preparations were generated according to published protocols (Klein, 2002). Briefly, 1mg of human A β ₁₋₄₂ (California peptide, catalog #641-15) was monomerized in 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP) and 0.45mg was aliquoted to non-siliconized microcentrifuge tubes. The HFIP was allowed to evaporate overnight in a hood at room temperature. If any HFIP remained, it was removed in a speed-vac for 10 minutes. A 5mM A β stock was then prepared by adding 20 μ l of anhydrous DMSO (Hybri-Max, Sigma) to 0.45mg of monomerized peptide film. Then, 980 μ l of Ham's F12 medium (BioSource, Inc) was added to create a 100 μ M oligomer solution. The resulting solution was incubated at 4°C for 24 hr.

Following incubation, the oligomer solution was centrifuged at 14,000 x g for 10 minutes at 4°C. The resulting supernatant was carefully recovered and used as the 100µM oligomer solution for cell toxicity studies.

Rat PC12 cells (ATCC) were plated at 20,000 cells/well in collagen-coated 96 well plates in F12K media (1% Horse serum, 1% Pen/Strep) and allowed to adhere overnight at 37°C and 5%CO₂. Media was refreshed with F12K just before the assay commenced. All Aβ antibodies (C700, C701, C705-707) and a commercially available mouse anti- Aβ antibody, 6E10, (Signet, catalog #9320-05) were diluted to 5.6µg/10µl in sterile water. Then, 5µM of Aβ₁₋₄₂ oligomers were pre-incubated with each of the antibodies for 2 hours at 4°C. Then, the oligomer and antibody combinations were added to the cells and incubated for 24 hr at 37°C. In this experiment, 5% ethanol was used as a positive control for cell toxicity. Cell viability was assessed by adding 10µl of MTT reagent (Roche, #1-465-007) to each well and allowed to incubate for 4hrs. Viable cells will reduce the MTT reagent to a formazan salt crystal. The crystals are solubilized overnight in the supplied buffer (Roche) and then read on a spectrophotometer at 550nm-690nm.

Results

The ability of the Aβ mAbs to inhibit Aβ₄₂ oligomer toxicity was tested using the rat PC12 cell line. Toxicity was measured using an MTT assay that determines cellular proliferation and viability. The MTT assay also represents a measure of cellular mitochondrial function since mitochondrial dehydrogenase activity is required to reduce the MTT dye to a formazan salt crystal, read on a spectrophotometer. There is typically a 40-50% decrease in MTT reduction following Aβ₄₂ oligomer exposure, as shown in Figure 48 upon comparison of Vehicle treated PC12 cells to those treated with 5 µM Aβ oligomers. The anti-human Aβ antibodies were tested for their ability to prevent of Aβ₄₂ oligomer toxicity. Aβ oligomers were pre-incubated with anti-human Aβ antibodies before they were exposed to the neuron-like PC12 cells.

Cellular exposure to 5µM oligomers alone resulted in a 27.3% decrease in cell viability compared to the vehicle control. All anti-human Aβ antibodies were able to impart neuroprotection on the PC12 cells following a 2 hr pre-incubation with the oligomers. Both C705 and C706 completely prevented Aβ₄₂ oligomer-induced toxicity in PC12 cells. The commercially available mouse monoclonal Ab antibody, 6E10, did not protect the cells at the concentration tested (560µg/ml).

It will be clear that the invention can be practiced otherwise than as particularly described in the foregoing description and examples.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

5

Table 1.

| SEQ ID NO | | | AA NO | regions | | | | | | |
|--------------|--------------------------------------|--------|----------|---------|------|-------|------|-------|------|--------|
| | | | | FR1 | CDR1 | FR2 | CDR2 | FR3 | CDR3 | FR4 |
| 1 | heavy chain variable region | Vh1 | 125 | 1-31 | 32 | 33-46 | 47 | 48-79 | 80 | 81-125 |
| 2 | | Vh2 | 97 | 1-30 | 31 | 32-45 | 46 | 47-78 | 79 | 80-97 |
| 3 | | Vh3a | 102 | 1-30 | 31 | 32-45 | 46 | 47-78 | 79 | 80-102 |
| 4 | | Vh3b | 102 | 1-30 | 31 | 32-45 | 46 | 47-78 | 79 | 80-102 |
| 5 | | Vh3c | 94 | 1-30 | 31 | 32-45 | 46 | 47-78 | 79 | 80-94 |
| 6 | | Vh4 | 106 | 1-30 | 31 | 32-45 | 46 | 47-78 | 79 | 80-106 |
| 7 | | Vh5 | 97 | 1-30 | 31 | 32-45 | 46 | 47-78 | 79 | 80-97 |
| 8 | | Vh6 | 91 | 1-30 | 31 | 32-45 | 46 | 47-78 | 79 | 80-91 |
| 9 | | Vh7 | 91 | 1-30 | 31 | 32-45 | 46 | 47-78 | 79 | 80-91 |
| 10 | light chain variable region | κ1_4 | 73 | 1-23 | 24 | 25-39 | 40 | 41-72 | 73 | |
| 11 | | κ2 | 73 | 1-23 | 24 | 25-39 | 40 | 41-72 | 73 | |
| 12 | | κ3 | 73 | 1-23 | 24 | 25-39 | 40 | 41-72 | 73 | |
| 13 | | κ5 | 73 | 1-23 | 24 | 25-39 | 40 | 41-72 | 73 | |
| 14 | | κ new1 | 67 | 1-17 | 18 | 19-33 | 34 | 35-66 | 67 | |
| 15 | | κ new2 | 65 | 1-15 | 16 | 17-31 | 32 | 33-64 | 65 | |
| 16 | | λ1a | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 17 | | λ1b | 73 | 1-23 | 24 | 25-39 | 40 | 41-72 | 73 | |
| 18 | | λ1c | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 19 | | λ3a | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 20 | | λ3b | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 21 | | λ3c | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 22 | | λ3e | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 23 | | λ4a | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 24 | | λ4b | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 25 | | λ5 | 75 | 1-22 | 23 | 24-39 | 40 | 41-74 | 75 | |
| 26 | | λ6 | 74 | 1-22 | 23 | 24-38 | 39 | 40-73 | 74 | |
| 27 | | λ7 | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 28 | | λ8 | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 29 | | λ9 | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |
| 30 | | λ10 | 72 | 1-22 | 23 | 24-38 | 39 | 40-71 | 72 | |

| SEQ ID NO | | | AA NO | regions | | | | | | | |
|--------------|--------------------------------------|------|----------|---------|---------|---------|---------|---------|---------|---------|---------|
| | | | | CH1 | hinge1 | hinge2 | hinge3 | hinge4 | CH2 | CH3 | CH4 |
| | | | | | | | | | | | |
| 31 | heavy chain constant region | IgA1 | 354 | 1-102 | 103-122 | | | | 123-222 | 223-354 | |
| 32 | | IgA2 | 340 | 1-102 | 103-108 | | | | 109-209 | 210-340 | |
| 33 | | IgD | 384 | 1-101 | 102-135 | 136-159 | | | 160-267 | 268-384 | |
| 34 | | IgE | 497 | 1-103 | | | | | 104-210 | 211-318 | 319-497 |
| 35 | | IgG1 | 339 | 1-98 | 99-113 | | | | 114-223 | 224-339 | |
| 36 | | IgG2 | 326 | 1-98 | 99-110 | | | | 111-219 | 220-326 | |
| 37 | | IgG3 | 377 | 1-98 | 99-115 | 116-130 | 131-145 | 146-160 | 161-270 | 271-377 | |
| 38 | | IgG4 | 327 | 1-98 | 99-110 | | | | 111-220 | 221-327 | |
| 39 | | IgM | 476 | 1-104 | | | | | 105-217 | 218-323 | 324-476 |
| 40 | light chain constant region | Igκc | 107 | | | | | | | | |
| 41 | | Igλc | 107 | | | | | | | | |

SEQUENCE LISTING

SEQ ID NO:1
 QVQLLVQSGA EVKKPGASVK VSCKASGYTF TXWVRQAPGQ GLEWMGXRV T MTRDTSTSTA
 5 60
 YMELSSLRSE DTAVYYCARX WQQTGLVTVS SGSTKGPSVF PLAPSSKSTS GGTAALGCLV
 120
 KDYFP
 125

10

SEQ ID NO:2
 QITLKESGPA LVKPTQTLTL TCTFSGFSL S XWIRQPPGKA LEWLAXRLTI SKDTSKNQVV
 60
 LTMNMDPVD TATYYCARXW GQGTGLVTVSS GSASAPS
 15 97

SEQ ID NO:3
 EVQLVESGGG LVQPGGSLRL SCAASGFTFS XWVRQAPGKG LEWVSXRFTI SRDNSKNTLY
 60
 20 LQMNSLRAED TAVYYCARXW GQGQGTGLVTV SSGSTKAPSV FP
 102

SEQ ID NO:4
 EVQLVESGGG LVKPGGSLRL SCAASGFTFS XWVRQAPGKG LEWVGXRFTI SRDDSKNTLY
 25 60
 LQMNSLKTED TAVYYCTTXW GQGTGLVTVSS ASTKGPSVFP LA
 102

SEQ ID NO:5
 30 EVQLVESGGG LVQPGRSLRL SCTASGFTFG XWVRQAPGKG LEWVGXRFTI SRDDSKSIAY
 60
 LQMNSLKTED TAVYYCTRXV TVSSGSTKGP SVLP
 94

35

SEQ ID NO:6
 QVQLQESGPG LVKPSETLSL TCTVSGGSIS XWIRQPPGKG LEWIGXRVTI SVDTSKNQFS
 60
 LKLSSVTAAD TAVYYCARXW GQGTGLVTVS SAPTKAPDVF PIISGC
 106

40

SEQ ID NO:7
 EVQLVQSGAE VKKPGESLKI SCKGSGYSFT XWVRQMPGKG LEWMGXQVTI SADKSISTAY
 60
 LQWSSLKASD TAVYYCARXW GQGTGLVTVSS ASTKGPS
 45 97
 CEN5021 NP

SEQ ID NO:8
 QVQLQQSGPG LVKPSQTLST TCAISGDSVS XWIRQSPSRG LEWLGXRITI NPDTSKNQFS
 60
 5 LQLNSVTPED TAVYYCARXW GQGTLVTVSS G
 91

SEQ ID NO:9
 QVQLVQSGSE LKKPGASVKV SCKASGYTFT XWVRQAPGQG LEWMGXRFVF SLDTSVSTAY
 10 60
 LQISSLKAED TAVYYCARXW GQGTLVTVSS S
 91

SEQ ID NO:10
 15 DIQMTQSPSS LSASVGDRVT ITCXWYQQKP GKAPKLLIYX GVPSRFSGSG SGTDFTLTIS
 60
 SLQPEDFATY YCX
 73

20 SEQ ID NO:11
 DIVMTQSPPLS LPVTPGQPAS ISCXWYLQKP GQSPQLLIYX GVPDRFSGSG SGTDFTLKIS
 60
 RVEAEDVGVI YCX
 73

25 SEQ ID NO:12
 EIVLTQSPGT LSLSPGERAT LSCXWYQQKP GQAPRLLIYX GIPDRFSGSG SGTDFTLTIS
 60
 RLEPEDFAVY YCX
 30 73

SEQ ID NO:13
 ETTLTQSPAF MSATPGDKVN ISCXWYQQKP GEAAIFIIQX GIPPRFSGSG YGTDFTLTIN
 60
 35 NIESEDAAYY FCX
 73

SEQ ID NO:14
 EIVMTQSPVN LSMSAGEXWY QQKPGQAPRL FIYXGISARF SGSGSGTDFT LTITSLQSED
 40 60
 FAVYYCX
 67

SEQ ID NO:15
 CEN5021 NP

ELTQSPGTLS LSPGEXWYQH KPGQAPRLVI HXGISDRFSG SGSGTDFTLT ITRLEPEDFA
 60
 LYYCX
 65
 5 SEQ ID NO:16
 QSVLTQPPSA SGTPGQRVTI SCXWYQQLPG TAPKLLIYXG VPDRFSGSKS GTSASLAISG
 60
 LQSEDEADYY CX
 10 72

 SEQ ID NO:17
 AQSVLTQPPS VSAAPGQKVT ISXWYQQLP GTAPKLLIYX GIPDRFSGSK SGTSATLGIT
 60
 15 GLQTGDEADY YCX
 73

 SEQ ID NO:18
 QSALTQPASV SGSPGQSITI SCXWYQQHPG KAPKLMIYXG VSNRFSGSKS GNTASLTISG
 20 60
 LQAEDEADYY CX
 72

 SEQ ID NO:19
 25 SYELTQPPSV SVSPGQTARI TCXWYQQKPG QAPVLVIYXG IPERFSGSSS GTTVTLTISG
 60
 VQAEDEADYY CX
 72

 30 SEQ ID NO:20
 SYVLTQPPSV SVAPGQTARI TCXWYQQKPG QAPVLVVYXG IPERFSGSNS GNTATLTISR
 60
 VEAGDEADYY CX
 72
 35
 SEQ ID NO:21
 SYELTQPPSV SVSPGQTASI TCXWYQQKPG QSPVLVIYXG IPERFSGSNS GNTATLTISG
 60
 TQAMDEADYY CX
 40 72

 SEQ ID NO:22
 SSELTDPAV SVALGQTVRI TCXWYQQKPG QAPVLVIYXG IPDRFSGSSS GNTASLTITG
 60

AQAEDEADYY CX
72

SEQ ID NO:23

5 QLVLTQSPSA SASLGASVKL TCXWHQQQPE KGPRYLMKXG IPDRFSGSSS GAERYLTISS
60
LQSEDEADYY CX
72

10 SEQ ID NO:24

QPVLTQSSSA SASLGSSVKL TCXWHQQQPG KAPRYLMKXG VPDRFSGSSS GADRYLTISN
60
LQSEDEADYY CX
72

15

SEQ ID NO:25

QAVLTQPSSL SASPGASASL TCXWYQQKPG SPPQYLLRYX GVPSRFSGSK DASANAGILL
60
ISGLQSEDEA DYICX

20 75

SEQ ID NO:26

NFMLTQPHSV SESPCKTVTI SCXWYQQRPG SAPTTVIYXG VPDRFSGSID SSSNSASLTI
60

25 SGLKTEDEAD YYCX
74

SEQ ID NO:27

30 QTVVTQEPSL TVSPGGTVTL TCXWFQQKPG QAPRALIYXW TPARFSGSLL GGKAALTLSG
60
VQPEDEAEYY CX
72

SEQ ID NO:28

35 QTVVTQEPSF SVSPGGTVTL TCXWYQQTPG QAPRTLIIYXG VPDRFSGSIL GNKAALTITG
60
AQADESDYY CX
72

40 SEQ ID NO:29

QPVLTQPPSA SASLGASVTL TCXWYQQRPG KGPRFVMRXG IPDRFSVLGS GLNRYLTIKN
60
IQEEDSDYH CX
72

SEQ ID NO:30

QAGLTQPPSV SKGLRQTATL TCXWLQQHQG HPPKLLSYXG ISERFSASRS GNTASLTITG
60
5 LQPEDEADYY CX
72

SEQ ID NO:31

ASPTSPKVFP LSLCSTQPDG NVVIACLVQG FFPQEPLSVT WSESGQGVTA RNFPSPQDAS
10 60
GDLYTTSSQL TLPATQCLAG KSVTCHVKHY TNPSQDVTVP CPVPSTPPTP SPSTPPTPSP
120
SCCHPRLSLH RPALEDLLL SEANLTCTLT GLRDASGVTF TWTPSSGKSA VQGPPERDLC
180
15 GCYSVSSVLP GCAEPWNHGK TFTCTAAYPE SKTPLTATLS KSGNTFRPEV HLLPPPSZEE
240
LALNELVTLT CLARGFSPKD VLVRWLQGSQ ELPREKYLTV ASRQEPSQGT TTFAVTSILR
300
VAAEDWKKGD TFSCMVGHEA LPLAFTQKTI DRLAGKPTHV NVSVVMAEVD GTCY
20 354

SEQ ID NO:32

ASPTSPKVFP LSLDSTPQDG NVVVIACLVQG FFPQEPLSVT WSESGQNVTA RNFPSPQDAS
60
25 GDLYTTSSQL TLPATQCPDG KSVTCHVKHY TNPSQDVTVP CPVPPPPPC HPRLSLHRPA
120
LEDLLLSEA NLTCTLTGLR DASGATFTWT PSSGKSAVQG PPERDLCGCY SVSSVLPGCA
180
QPWNHGETFT CTAHPPELKT PLTANITKSG NTFRPEVHLL PPPSEELALN ELVTLTCLAR
30 240
GFSPKDVLR WLQGSQELPR EKYLTVASRQ EPSQGTTFV VTSILRVAAE DWKKGDTFSC
300
MVGHEALPLA FTQKTIDRLA GKPTHVNVSV VMAEVDGTCY
340

35

SEQ ID NO:33

APTKAPDVFP IISGCRHPKD NSPVVLACLI TGYHPTSVTV TWYMGTSQSP QRTFPEIQR
60
DSYYMTSSQL STPLQWRQG EYKCVVQHTA SKSKKEIFRW PESPKAQASS VPTAQQAEG
40 120
SLAKATTAPA TTRNTGRGGE EKKKEKEKEE QEERETKTPE CPSHTQPLGV YLLTPAVQDL
180
WLRDKATFTC FVVGSDLKDA HLTWEVAGKV PTGGVEEGLL ERHSNGSQSQ HSRLTLPRSL
240
45 WNAGTSVTCT LNHPSLPPQR LMALREPAAQ APVKLSLNL ASSDPPEAAS WLLCEVSGFS
CEN5021 NP

300
 PPNILLMWLE DQREVNTSGF APARPPPQPR STTFWAWSVL RVPAPPSPQP ATYTCVVSHE
 360
 DSRTLLNASR SLEVSIVTDH GPMK
 5 384

SEQ ID NO:34
 ASTQSPSVFP LTRCCKNIPS NATSVTLGCL ATGYFPEPVM VTWDTGSLNG TTMTLPATTL
 60
 10 TLSGHYATIS LLTVSGAWAK QMFTCRVAHT PSSTDWVDNK TFSVCSRDFE PPTVKILQSS
 120
 CDGGGHFPPT IQLLCLVSGY TPGTINITWL EDGQVMDVDL STASTQEGER LASTQSELTL
 180
 SQKHWLSDRY YTCQVTYQGH TFEDSTKKCA DSNPRGVSAY LSRPSPFDLF IRKSPTITCL
 15 240
 VVDLAPSKGT VNLTWSRASG KPVNHSTRKE EKQRNGTLTV TSTLPVGTRD WIEGETYQCR
 300
 VTHPHLPRAL MRSTTKTSGP VGPRAAPEVY AFATPEWPGS RDKRTLACLI QNFMPEDISV
 360
 20 QWLHNEVQLP DARHSTTQPR KTKGSGFFVF SRLEVTRAEW EQKDEFICRA VHEAASPSQT
 420
 VQRAVSVNPG KDVCVEEAEG EAPWTWTGLC IFAALFLLSV SYSAALTLLM VQRFLSATRQ
 480
 GRPQTSLDYT NVLQPHA
 25 497

SEQ ID NO:35
 ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS
 60
 30 GLYSLSSVVT VPSSSLGTQT YICNVNHKPS NTKVDKKVEP KSCDKTHTCP PCPAPELLGG
 120
 PSVFLFPKP KDTLMISRTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN
 180
 STYRVVSVLT VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSRDE
 15 240
 LTKNQVSLTC LVKGFYPSDI AVEWESBNGQ PENNYKTPP VLDSGDSFFL YSKLTVDKSR
 300
 WQQGNVFSCS VMHEALHNHY TQKSLSLSPG KTHTCPPCP
 339
 40

SEQ ID NO:36
 ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS
 60
 45 GLYSLSSVVT VPSSNFGTQT YTCNVNHHKPS NTKVDKTVR KCCVECPPCP APPVAGPSVF
 120
 CEN5021 NP

LFPPKPKDTL MISRTPEVTC VVVDVSHEDP EVQFNWYVDG VEVHNAKTKP REEQFNSTFR
 180
 VVSVLTVVHQ DWLNGKEYKC KVSNGKLPAP IEKTISKTKG QPREPQVYTL PPSREEMTKN
 240
 5 QVSLTCLVKG FYPSDIAVEW ESNGQPENNY KTTTPMLDSD GSFFLYSKLT VDKSRWQQGN
 300
 VFSCSVMHEA LHNHYTQKSL SLSPGK
 326

10 SEQ ID NO:37
 ASTKGPSVFP LAPCSRSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS
 60
 GLYSLSSVVT VPSSSLGTQT YTCNVNHKPS NTKVDKRVEL KTPLGDTTHT CPRCPEPKSC
 120
 15 DTPPPCPRCP EPKSCDTPPP CPRCPEPKSC DTPPPCPRCP APELLGGPSV FLFPPKPKDT
 180
 LMISRTPEVT CVVVDVSHED PEVQFKWYVD GVEVHNAKTK PREEQYNSTF RVVSVLTVLH
 240
 QDWLNGKEYK CKVSNKALPA PIEKTISKTK GQPREPQVYT LPPSREEMTK NQVSLTCLVK
 20 300
 GFYPSDIAVE WESSGQPENN YNTTPMLDS DGSFFLYSKL TVDKSRWQQG NIFSCSVMHE
 360
 ALHNRFTQKS LSLSPGK
 377

25 SEQ ID NO:38
 ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS
 60
 GLYSLSSVVT VPSSSLGTKT YTCNVNHNKPS NTKVDKRVES KYGPPCPSCP APEFLGGPSV
 30 120
 FLFPPKPKDT LMISRTPEVT CVVVDVSQED PEVQFNWYVD GVEVHNAKTK PREEQFNSTY
 180
 RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS SIEKTISKAK GQPREPQVYT LPPSQEEMTK
 240
 35 NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSRL TVDKSRWQEG
 300
 NVFSCSVMHE ALHNHYTQKS LSLSLGK
 327

40 SEQ ID NO:39
 GSASAPTLFP LVSCENSPSD TSSVAVGCLA QDFLPDSITF SWKYKNNSDI SSTRGFPSVL
 60
 RGGKYAATSQ VLLPSKDVMQ GTDEHVCKV QHPNGNKEKN VPLPVIAELP PKVSVFVPPR
 120
 45 DGFFGNPRSK SKLICQATGF SPRQIQVSWL REGKQVGSGV TTDQVQAEAK ESGPTTYKVT
 CEN5021 NP

| | | |
|----|--|-----|
| | 180 | |
| | STLTIKESDW LSQSMFTCRV DHRGLTFQQN ASSMCVPDQD TAIRVFAIPP SFASIFLTKS | |
| | 240 | |
| | TKLTCLVTDL TTYDSVTISW TRQNGEAVKT HTNISESHPN ATFSAVGEAS ICEDDWNNGE | |
| 5 | 300 | |
| | RFTCTVTHTD LPSPLKQTIS RPKGVALHRP DVYLLPPARE QNLNRESATI TCLVTGFSPA | |
| | 360 | |
| | DVVFVQWQMR GQPLSPEKYV TSAPMPEPQA PGRYFAHSIL TVSEEEWNTG ETYTCVVAHE | |
| | 420 | |
| 10 | ALPNRV TERT VDKSTGKPTS ADEEGFENLW ATASTFIVLY NVSLVMSDTA GTCYVK | |
| | 476 | |
| | SEQ ID NO:40 | |
| | RTVAAPSVFI FPPSDEQLKS GTASVVCLLN NFYPREKAVQ WKVDNALQSG NSQESVTEQD | |
| 15 | 60 | |
| | SKDSTYSLSS TLTLKADYE KHKVYACEVT HQGLSSPVTK SFNRGEC | |
| | 107 | |
| | SEQ ID NO:41 | |
| 20 | GQPKAAPSVT LFPPSSEELQ ANKATLVCLI SDFYPGAVTV AWKADSSPVK AGVETTTPSK | |
| | 60 | |
| | QSNKYAASS YLSLTPEQWK SHRKSYSCQV THEGSTVEKT VAPTECS | |
| | 107 | |
| 25 | SEQ ID NO:42 | |
| | DHYVH | 5 |
| | SEQ ID NO:43 | |
| | WIAPKNGYSE SAPKFQG | 17 |
| 30 | | |
| | SEQ ID NO:44 | |
| | GFYDSSLY | 8 |
| | SEQ ID NO:45 | |
| 35 | KSGQSL LARD GKTYLS | 16 |
| | SEQ ID NO:46 | |
| | LVS KLDS | 7 |
| 40 | SEQ ID NO:47 | |
| | WQGT HFPRT | 9 |
| | SEQ ID NO:48 | |
| | MKCSWVIFFL MAVVIGINSE GQLQQSGAEL VRSGASLKLS CTASGFNIKD HYVHWVRQRP | 60 |
| 45 | EQGLDWIGWI APKNGYSESA PKFQ GKASMT ADTSSNTVYL QLSSLTSEDV AVYYCFAGFY | 120 |
| | DSSLYWGQGT TLT VSS | 136 |
| | CEN5021 NP | 100 |

| | | |
|----|---|-----|
| | SEQ ID NO:49 | |
| | MMSPAQFLFL LVLWIRETNG DVVMTQTPLT LAVTIGQPAS ISCKSGQSL L ARDGKTYLSW | 60 |
| | LLQRPQGSPK RLIYLVSKLD SGVPDRFSGS GSGTDFTLKI NRVEAEDLGV YYCWQGTHFP | 120 |
| 5 | RTFGGGTNLE IKR | 133 |
| | SEQ ID NO:50 | |
| | DAEFRHDSGY EVHHQKLVFF AEDVGSNKGA IIGLMVGGVV IA | 42 |
| 10 | SEQ ID NO:51 | |
| | atgaaatgca gctgggtcat cttcttcctg atggcagtgg tcataggaat caattcagag | 60 |
| | ggtcagctgc agcagtctgg ggcagaactt gtgagggtcag gggcctcact caagttgtcc | 120 |
| | tgcacagctt ctggcttcaa tattaaagac cactatgtac actgggtgag gcagaggcct | 180 |
| | gaacagggcc tggactggat tggatggatt gctccgaaga atgggttatag tgaatctgcc | 240 |
| 15 | ccgaaattcc agggcaaggc cagtatgact gcagacacat cctccaacac agtctacctg | 300 |
| | cagctcagca gcctgacatc tgaggacact gccgtctatt actgttttgc agggttttac | 360 |
| | gatagtagcc tctactgggg ccagggcacc actctcacag tctcttca | 408 |
| | SEQ ID NO:52 | |
| 20 | atgatgagtc ctgcccagtt cctgtttctg ttagtgctct ggattcggga aaccaacggt | 60 |
| | gacgttgtaa tgaccagac tccactcact ttggcggtta ccattggaca accagcctcc | 120 |
| | atctcttgca agtcaggta gagcctctta gcaagagatg gaaagacata tttgagttgg | 180 |
| | ttattacaga ggccaggcca gtctccaaag cgcctaactc atctggtgtc taaactggac | 240 |
| | tctggagtc ctgacaggtt ctctggcagt ggatcaggga cagatttcac actgaaaatc | 300 |
| 25 | aacagagtgg aggctgagga tttgggagtt tattattgct ggcaaggtag acattttcct | 360 |
| | cggacgttcg gtggaggcac caacctggaa atcaaacgg | 399 |
| | SEQ ID NO:53 | |
| 30 | TSGMGVS | 7 |
| | SEQ ID NO:54 | |
| | HIYWDDDKRY NPSLKS | 16 |
| | SEQ ID NO:55 | |
| 35 | SSGSIVIATG FAY | 13 |
| | SEQ ID NO:56 | |
| | RSSQSLVHSN GNTYLE | 16 |
| 40 | SEQ ID NO:57 | |
| | KVSNRFS | 7 |
| | SEQ ID NO:58 | |
| 45 | FQGSRVPLT | 9 |
| | SEQ ID NO:59 | |
| | MDRLTSSFL L LIVPAYVLSQ VTLKESGPGI LQPSQTL SLT CSFSGFSLST SGMGVSWIRQ | 60 |
| | CEN5021 NP | 101 |

| | | |
|----|--|-----|
| | PSGKGLEWLA HIYWDDDKRY NPSLKSRLTI SKDTSRNQVF LKITSVDTTD TATYYCTRSS | 120 |
| | GSIVIATGFA YWGQGLVTV SA | 142 |
| | SEQ ID NO:60 | |
| 5 | MKLPVRLVL MFWIPGSSSD VMMTQTPLSL PVSLGDQASI SCRSSQSLVH SNGNTYLEWY | 60 |
| | MQKPGQSPML LIYKVSNRFS GVPDRFSGSG SGTDFTLKIS SVEAEDLGVF YCFQGSRVPL | 120 |
| | TFGAGTKLEL KR | 132 |
| | SEQ ID NO:61 | |
| 10 | atggacaggc ttacttcctc attcctgctg ctgattgtcc ctgcatatgt cctttcccag | 60 |
| | gttactctga aagagtctgg ccctgggata ttgcagccct ccagaccct cagtctgact | 120 |
| | tgttctttct ctgggttttc actgagcact tctggtatgg gtgtgagctg gattcgtcag | 180 |
| | ccttcaggaa agggctctgga gtggctggca cacatttact gggatgatga caaacgatat | 240 |
| | aatccatccc tgaagagccg gctcacaatc tccaaggata cttccagaaa ccaggtattc | 300 |
| 15 | ctcaagatca ccagtgtgga cactacagat actgccacat actactgtac tcgaagttcc | 360 |
| | ggatctattg tgattgcgac ggggtttgct tactggggcc aagggactct ggtcactgtc | 420 |
| | tctgca | 426 |
| | SEQ ID NO:62 | |
| 20 | atgaagttgc ctggttaggct gttggtgctg atgttctgga ttcctggttc cagcagtgat | 60 |
| | gttatgatga cccaaactcc actctccctg cctgtcagtc ttggagatca agcctccatc | 120 |
| | tcttgcatat ctagtccagag tcttgtagat agtaatggaa acacctattt agaatggat | 180 |
| | atgcagaaac caggccagtc tccaatgctc ctgatctaca aagtttccaa ccgattttct | 240 |
| | ggggtcccag acaggttcag tggcagtgga tcaggacag atttcacact caagatcagc | 300 |
| 25 | agcgtggagg ctgaggatct gggagttttt tactgctttc aagggttcacg tgttccgctc | 360 |
| | acgttcggtg ctgggaccaa gctggagctg aaacgg | 396 |
| | SEQ ID NO:63 | |
| 30 | TSWIE | 5 |
| | SEQ ID NO:64 | |
| | EVLPGSGKSN HNANFKG | 17 |
| | SEQ ID NO:65 | |
| 35 | EGSNNNALAY | 10 |
| | SEQ ID NO:66 | |
| | SASSSVSYM | 10 |
| 40 | SEQ ID NO:67 | |
| | DSSRLAS | 7 |
| | SEQ ID NO:68 | |
| 45 | QNWRSSPT | 8 |
| | SEQ ID NO:69 | |
| | MEWTWVFLFL LSVTAGVHSQ VQLQQSGPEL MKPGASVKIS CKATGYTFST SWIEWIKQRP | 60 |

| | | |
|----|--|-----|
| | GHGLEWIGEV LPGSGKSNHN ANFKGRATFT ADTASNTAYM QLSSLTSEDS AVYYCAREGS | 120 |
| | NNNALAYWGQ GTLVTVSA | 138 |
| | SEQ ID NO:70 | |
| 5 | MDFQVQIFSF LLISASVIIS RGQIVLTQSP AIMSASPGEK VTMTCSASSS VSYMHWYQQK | 60 |
| | SGTSPKRWIY DSSRLASGVP SRFSGGSGT SYSPTISNME AEDAATYFCQ NWRSSPTFGA | 120 |
| | GTKLELKR | 128 |
| | SEQ ID NO:71 | |
| 10 | atggaatgga cctgggtctt tctcttcttc ctgtcagtaa ctgcaggtgt ccactcccag | 60 |
| | gttcaactgc agcagctctgg acctgaactg atgaagcctg gggcctcagt gaagatatcc | 120 |
| | tgcaaggcta ctggctacac attcagtacc tcctggatag agtggataaa gcagaggcct | 180 |
| | ggacatggcc ttgagtggat tggagaggtc ttacctggaa gcggtaaagag taaccacaat | 240 |
| | gcgaacttta agggcagggc cacatttact gcagatacag cctccaacac agcctacatg | 300 |
| 15 | cagctcagca gcctgacatc tgaggactct gccgtctatt attgtgcaag agaggggagt | 360 |
| | aataacaacg ctttggctta ctggggccaa gggactctgg tcaactgtctc tgca | 414 |
| | SEQ ID NO:72 | |
| | atggattttc aagtgcagat tttcagcttc ctgctaataca gtgcctcagt cataatatcc | 60 |
| 20 | agaggacaaa ttgtgctcac ccagtctcca gcaatcatgt ctgcttctcc aggggagaag | 120 |
| | gtcaccatga cctgcagtgc cagctcaagt gtgagttaca tgcaactggta ccaacagaag | 180 |
| | tcaggcacct cccccaaaag atggatttat gacagttcca gactggcttc tggagtccct | 240 |
| | tctcgcttca gtggcggtgg gtctgggacc tcttactctc ccacaatcag caacatggag | 300 |
| | gctgaagatg ctgccacgta tttctgccag aactggcgta gtagccccac gttcgggtgct | 360 |
| 25 | gggaccaagc tggagctgaa acgg | 384 |
| | SEQ ID NO:73 | |
| | EYIMS | 5 |
| 30 | SEQ ID NO:74 | |
| | SINPNTGGSR YNQKFKG | 17 |
| | SEQ ID NO:75 | |
| | GDFDY | 5 |
| 35 | SEQ ID NO:76 | |
| | RSSKNLLHSN GITYLY | 16 |
| | SEQ ID NO:77 | |
| 40 | RVSNLAS | 7 |
| | SEQ ID NO:78 | |
| | AQLLELPFT | 9 |
| 45 | SEQ ID NO:79 | |
| | MGWSWIFLFL LSGTAGVLSE VQLQQSGPDL VKPGASVKTS CKTSGYSFTE YIMSWVRQSH | 60 |
| | GKSLEWIGSI NPNTGGSRYN QKFKGKATLT VDKSSSTAYM EFRSLTSEDS AVYYCARGDF | 120 |
| | CEN5021 NP | |

| | | |
|----|---|-----|
| | DYWGQGTTLT VSS | 133 |
| | SEQ ID NO:80 | |
| | MRFSAQLLGL LVLWIPGSTA DIVMTQAAFS NPVTLGTSAS ISCRSSKNLL HSNGITYLYW | 60 |
| 5 | YLQRPQGSPQ LLISRVSNLA SGVPNRFSGS ESGTDFTLRI SRVEAEDVGV YYCAQLLELP | 120 |
| | FTFGSGTKLE IKR | 133 |
| | SEQ ID NO:81 | |
| | atgggatgga gctggatctt tctctttctc ctgtcaggaa ctgcaggtgt cctctctgag | 60 |
| 10 | gtccagctgc aacaatctgg acctgacctg gtgaagcctg gggcttcagt gaagacatcc | 120 |
| | tgcaagactt ctggatactc attcactgaa tacatcatga gctgggtgag gcagagccat | 180 |
| | ggaaagagcc ttgagtggat tggaagtatt aatcctaaca ctggtggtag tagatacaac | 240 |
| | cagaaattca agggcaaggc cacgttgact gtagataagt cctccagcac agcctacatg | 300 |
| | gagtttcgca gcctgacatc tgaggattct gcagtctatt actgtgcaag aggggacttt | 360 |
| 15 | gactactggg gcccaaggcac cactctcaca gtctcctca | 399 |
| | SEQ ID NO:82 | |
| | atgaggttct ctgctcagct tctggggctg cttgtgctct ggatccctgg atccactgca | 60 |
| | gatattgtga tgacgcaggc tgccttctcc aatccagtca ctcttggaac atcagcttcc | 120 |
| 20 | atctcctgca ggtctagtaa gaatctccta catagtaatg gcatcactta tttgtattgg | 180 |
| | tatctgcaga ggccaggcca gtctcctcag ctccctgatat ctcggtgtc caatctggcc | 240 |
| | tcaggagtcc caaacaggtt cagtggcagt gagtcaggaa ctgatttcac actgagaatc | 300 |
| | agcagagtgg aggctgagga tgtgggtgtt tattactgtg ctcaactgct agaactccca | 360 |
| | ttcacgttcg gctcggggac aaagtggaa ataaaaagg | 399 |
| 25 | | |
| | AEFRHDSGYEVH (SEQ ID NO:83); | |
| | EFRHDSGYEVHH (SEQ ID NO:84); | |
| 30 | IIGLMVGGVVIA (SEQ ID NO:85); | |
| | IIGLMVGGVVIA (SEQ ID NO:86); | |
| | IIGLMVGGVVI (SEQ ID NO:87); | |
| 35 | IIGLMVGGVV (SEQ ID NO:88); | |
| | IIGLMVGGV (SEQ ID NO:89); | |
| 40 | IIGLMVGG (SEQ ID NO:90); | |
| | LMVGGV (SEQ ID NO:91). | |
| | DAEFRHDSGYEVHHQ (SEQ ID NO:92); | |
| 45 | AEFRHDSGYEVHHQ(SEQ ID NO:93); | |
| | DAEFRHDSGYEVH(SEQ ID NO:94); | |
| | CEN5021 NP | 104 |

EFRHDSGYEVHH(SEQ ID NO:95);
 EFRHDSGYEVH(SEQ ID NO:96);
 5 DAEFRHDSGY(SEQ ID NO:97);
 DAEFRHDSG(SEQ ID NO:98);
 10 AEFRHDSG(SEQ ID NO:99);
 EFRHDSG (SEQ ID NO:100);
 EFRHDS (SEQ ID NO:101)
 15 DAEFRHDSGYEVHHQ (SEQ ID NO:102);
 AEFRHDSGYEVHHQ (SEQ ID NO:103);
 20 AEFRHDSGYEVHH (SEQ ID NO:104);
 AEFRHDSGYEVH (SEQ ID NO:105);
 DAEFRHDSGYE (SEQ ID NO:106);
 25 AEFRHDSGYE (SEQ ID NO:107);
 DAEFRHDSG (SEQ ID NO:108);
 30 AEFRHDSG (SEQ ID NO:109);
 DAEFRHDSG (SEQ ID NO:110);
 AEFRHD (SEQ ID NO:111)
 35 EFRHDSGYEVHHQKL (SEQ ID NO:112);
 FRHDSGYEVHHQKL (SEQ ID NO:113);
 40 EVHHQKLVFFAEDV (SEQ ID NO:114);
 YEVHHQKLVFFA (SEQ ID NO:115);
 VFFAEDVGSNKGA (SEQ ID NO:116);
 45 KLVFFAEDVGSN (SEQ ID NO:117);
 EDVGSNKGAIIG (SEQ ID NO:118);
 50 FFAEDVGSNKG (SEQ ID NO:119);
 CEN5021 NP

SNKGAIIGLMV (SEQ ID NO:120);
FAEDVGSNKG (SEQ ID NO:121);
5 KGAIIGLMVG (SEQ ID NO:122);
LVFFAEDVG (SEQ ID NO:123);
10 EDVGSNKGA (SEQ ID NO:124);
KLVFFAED (SEQ ID NO:125);
DVGSNKGA (SEQ ID NO:126);
15 EVHHQKL (SEQ ID NO:127);
QKLVFFA (SEQ ID NO:128);
20 RHDSGY (SEQ ID NO:129);
SGYEVH (SEQ ID NO:130);
GVVIAT (SEQ ID NO:131).
25